

Differential Equations Compatible with KZ Equations

G. Felder^{*}, Y. Markov^{*}, V. Tarasov^{**} and A. Varchenko[◇]

^{*} *Departement Mathematik, ETH-Zentrum,
8092 Zürich, Switzerland*

^{*,◇} *Department of Mathematics, University of North Carolina,
Chapel Hill, NC 27599 – 3250, USA*

^{**} *St. Petersburg Branch of Steklov Mathematical Institute
Fontanka 27, St. Petersburg, 191011, Russia*

January, 2000

1 Introduction

In the theory of the bispectral problem [DG], [W], one considers a commutative algebra A of differential operators $L(z, \partial/\partial z)$ acting on functions of one complex variable z . Such an algebra is called bispectral if there exists a non-trivial family $u(z, \mu)$ of common eigenfunctions depending on a spectral parameter μ

$$Lu(z, \mu) = f_L(\mu)u(z, \mu), \quad L \in A, \quad (1)$$

which is also a family of common eigenfunctions of a commutative algebra B of differential operators $\Lambda(\mu, \partial/\partial \mu)$ with respect to μ :

$$\Lambda u(z, \mu) = \theta_\Lambda(z)u(z, \mu), \quad \Lambda \in B. \quad (2)$$

J. Duistermaat and A. Grünbaum [DG] studied the case where A is the algebra of differential operators that commute with a Schrödinger operator $\frac{d^2}{dz^2} - V(z)$ with meromorphic potential $V(z)$. They give a complete classification of bispectral algebras arising in this way. In

^{*} *E-mail: felder@math.ethz.ch*

^{*} *E-mail: markov@math.unc.edu*

^{**} *E-mail: vt@pdmi.ras.ru*

[◇] *E-mail: av@math.unc.edu*

Supported in part by NSF grant DMS-9801582

particular they show that A is bispectral if $V(z)$ is a rational KdV potential (a rational function which stays rational under the flow of the Korteweg–de Vries equation). G. Wilson [W] classified bispectral algebras of rank one, i.e., such that the greatest common divisor of the orders of the differential operators in A is one. He showed that the maximal bispectral algebras of rank one are in one to one correspondence with conjugacy classes of pairs (Z, M) of square matrices so that $ZM - MZ + I$ has rank one. The bispectrality then follows from the existence of the involution $(Z, M) \mapsto (M^T, Z^T)$, which corresponds to exchanging z and μ .

The higher dimensional version of the bispectral problem, in which A consists of partial differential operators in $z \in \mathbb{C}^n$ is open. However O. Chalykh, M. Feigin and A. Veselov [CV],[CVF] constructed examples of algebras in higher dimensions which have the bispectral property (see Veselov’s contribution to [HK]). In these examples A consists of differential operators commuting with an n -particle Schrödinger operator with certain special rational potentials, including the Calogero–Moser ones. These potentials are in many respects the natural generalization of rational KdV potential associated to rank one algebras. In these examples, the Baker–Akhiezer function $u(z, \mu)$ is symmetric in the two arguments, thus $B = A$.

A good source of material on the bispectral problem is the volume [HK].

In this paper we study a class of examples of commutative algebras of partial differential operators acting on *vector-valued* functions with the bispectral property. This means that in (1), (2), u takes values in a vector space and $f_L(\mu)$, $\theta_\Lambda(z)$ are endomorphisms of the vector space. In our class of examples, the algebra A is generated by Knizhnik–Zamolodchikov differential operators. They are commuting first order differential operators associated to a complex simple Lie algebra \mathfrak{g} with a fixed non-degenerate invariant bilinear form and a non-zero complex parameter κ . They act on functions of n complex variables z_i taking values in the tensor product of n finite dimensional \mathfrak{g} -modules. The “dual” variable μ is in a Cartan subalgebra of \mathfrak{g} . The first set of equations (1) is then the set of generalized Knizhnik–Zamolodchikov equations

$$\left(\kappa \frac{\partial}{\partial z_i} - \sum_{j:j \neq i} \frac{\Omega^{(ij)}}{z_i - z_j} \right) u(z, \mu) = \mu^{(i)} u(z, \mu), \quad i = 1, \dots, n.$$

Here $\Omega \in \mathfrak{g} \otimes \mathfrak{g}$ is dual to the invariant bilinear form and $\Omega^{(ij)}$ acts as Ω on the i th and j th factors of the tensor product and as the identity on the other factors. Similarly $\mu^{(i)}$ is μ acting on the i th factor. It is well-known that these equations form a compatible system, i.e., they are the equations defining horizontal sections for a flat connection. For $\mu = 0$ they reduce to the classical Knizhnik–Zamolodchikov equations. The algebra B is generated by $\text{rank}(\mathfrak{g})$ first order partial differential operators in μ with rational coefficients. We call the corresponding equations (2) *dynamical differential equations*, and show that they form, together with the generalized Knizhnik–Zamolodchikov equations, a compatible system. We also give simultaneous solutions of both systems of equations in terms of hypergeometric integrals for a more general class of Lie algebras, which includes in particular all Kac–Moody Lie algebras.

In the case of $\mathfrak{g} = sl_2$, the algebra B is generated by one ordinary differential operator. In this case, the corresponding equations were first written and solved by H. Babujian and A. Kitaev [BK], who also related the equations to the Maxwell–Bloch system.

Our paper is organized as follows. In Sect. 2 we introduce the systems of Knizhnik–Zamolodchikov and dynamical differential equations for arbitrary simple Lie algebra and prove their compatibility. We then give formulae for hypergeometric solutions in Sect. 3 and give as an application a determinant formula. The fact that the hypergeometric integrals provide solutions is a consequence of a general theorem valid for a class of Lie algebras with generic Cartan matrix, introduced in [SV]. We introduce in Sect. 4 the Knizhnik–Zamolodchikov and dynamical differential equations in this more general context and explain in the next Section the results on complexes of hypergeometric differential forms from [SV]. In Sect. 6 we prove that the hypergeometric integrals for generic Lie algebras satisfies the dynamical differential equations. Finally in Sect. 7 we prove that our hypergeometric integrals are solutions of both systems of equations for any Kac–Moody Lie algebra. We also find a determinant formula, which implies a completeness result for solutions in the case of generic parameters.

2 Dynamical differential equations

2.1

Let \mathfrak{g} be a simple complex Lie algebra with an invariant bilinear form (\cdot, \cdot) and a root space decomposition $\mathfrak{g} = \mathfrak{h} \oplus (\oplus_{\alpha \in \Delta} \mathbb{C}e_\alpha)$. The root vectors e_α are normalized so that $(e_\alpha, e_{-\alpha}) = 1$. Then the quadratic Casimir element of $\mathfrak{g} \otimes \mathfrak{g}$ has the form $\Omega = \sum_s h_s \otimes h_s + \sum_{\alpha \in \Delta} e_\alpha \otimes e_{-\alpha}$, for any orthonormal basis (h_s) of the Cartan subalgebra \mathfrak{h} . We also fix a system of simple roots $\alpha_1, \dots, \alpha_r$.

Consider the Knizhnik–Zamolodchikov (KZ) equations with an additional parameter $\mu \in \mathfrak{h}$, for a function u on n variables taking values in a tensor product $V = V_1 \otimes \dots \otimes V_n$ of highest weight modules of \mathfrak{g} with corresponding highest weights $\Lambda_1, \dots, \Lambda_n$,

$$\kappa \frac{\partial u}{\partial z_i} = \mu^{(i)} u + \sum_{i \neq j} \frac{\Omega^{(ij)}}{z_i - z_j} u, \quad i = 1, \dots, n, \quad (3)$$

where κ is a complex parameter. We are interested in a differential equation for u with respect to μ which are compatible with KZ equations. If $\mu' \in \mathfrak{h}$, denote by $\partial_{\mu'}$ the partial derivative with respect to μ in the direction of μ'

Theorem 2.1 *The equations*

$$\kappa \partial_{\mu'} u = \sum_{i=1}^n z_i (\mu')^{(i)} u + \sum_{\alpha > 0} \frac{\langle \alpha, \mu' \rangle}{\langle \alpha, \mu \rangle} e_{-\alpha} e_\alpha u, \quad \mu' \in \mathfrak{h} \quad (4)$$

form together with the KZ equations (3), a compatible system of equations for a function $u(z, \mu)$ taking values in $V = V_1 \otimes \dots \otimes V_n$.

The equations (4) will be called *dynamical differential equations*.

Example. Let $\mathfrak{g} = sl_N = gl_N/\mathbb{C}$. View sl_N -modules as gl_N -modules by letting the center of gl_N act trivially. Denote by $E_{a,b} \in gl_N$ the matrix whose entries are zero except for a one at the intersection of the a th row with the b th column. The fundamental coweights $\varpi_a = (1 - a/N) \sum_{b \leq a} E_{b,b} - (a/N) \sum_{b > a} E_{b,b}$, $a = 1, \dots, N-1$ form a basis of the standard Cartan subalgebra of sl_N . Write $\mu = \sum_{a=1}^{N-1} \mu_a \varpi_a$. Then our equations may be written as

$$\begin{aligned} \kappa \frac{\partial u}{\partial z_i} &= \sum_{a=1}^{N-1} \mu_a \varpi_a^{(i)} u + \sum_{j:j \neq i} \sum_{a,b} \frac{E_{a,b}^{(i)} E_{b,a}^{(j)}}{z_i - z_j} u. \\ \kappa \frac{\partial u}{\partial \mu_a} &= \sum_{i=1}^n z_i \varpi_a^{(i)} u + \sum_{b,c:b \leq a < c} \sum_{i,j} \frac{E_{b,c}^{(i)} E_{c,b}^{(j)}}{\mu_b + \mu_{b+1} \cdots + \mu_{c-1}} u. \end{aligned}$$

2.2 Proof of Theorem 2.1

It is rather easy to verify that most terms of the compatibility equations vanish. The only non-trivial thing to check is that the operators $\sum_{\alpha > 0} \frac{\langle \alpha, \lambda \rangle}{\langle \alpha, \mu \rangle} e_{-\alpha} e_{\alpha}$ commute for different values of λ . The operators obtained by extending the sum to all roots differ from the sum over positive roots by an element of the Cartan subalgebra. Since the operators commute with the Cartan subalgebra, it is sufficient to prove the following Proposition.

Proposition 2.2 *Let for $\lambda, \mu \in \mathfrak{h}$, $T(\lambda, \mu) = \sum_{\alpha \in \Delta} \frac{\langle \alpha, \lambda \rangle}{\langle \alpha, \mu \rangle} e_{-\alpha} e_{\alpha}$. Then for all $\lambda, \mu \in \mathfrak{h}$,*

$$T(\lambda, \mu)T(\nu, \mu) = T(\nu, \mu)T(\lambda, \mu).$$

The proof is based on the following fact.

Lemma 2.3 *Let $\alpha, \beta \in \Delta$ with $\alpha \neq \pm\beta$, and let $S = S(\alpha, \beta)$ be the set of integers j such that $\beta + j\alpha \in \Delta$. Then $\sum_{j \in S} [e_{\alpha}, e_{\beta+j\alpha} e_{-\beta-j\alpha}] = 0$ and $\sum_{j \in S} [e_{-\alpha}, e_{\beta+j\alpha} e_{-\beta-j\alpha}] = 0$*

Proof. For roots γ, δ such that $\gamma + \delta$ is a root, let $N_{\gamma, \delta} = ([e_{\gamma}, e_{\delta}], e_{-\gamma-\delta})$, so that $[e_{\gamma}, e_{\delta}] = N_{\gamma, \delta} e_{\gamma+\delta}$. By considering the adjoint action on \mathfrak{g} of the sl_2 sub-algebra generated by $e_{\pm\alpha}$, we see that for $\beta \neq \pm\alpha$, S is a finite sequence of subsequent integers. We may thus assume that $S = \{0, \dots, k\}$ by replacing β by $\beta - j\alpha$ for some j if necessary.

We then have

$$\begin{aligned} & \sum_{j=0}^k [e_{\alpha}, e_{\beta+j\alpha} e_{-\beta-j\alpha}] \\ &= \sum_{j=0}^{k-1} N_{\alpha, \beta+j\alpha} e_{\beta+(j+1)\alpha} e_{-\beta-j\alpha} + \sum_{j=1}^k N_{\alpha, -\beta-j\alpha} e_{\beta+j\alpha} e_{-\beta-(j-1)\alpha} \\ &= \sum_{j=0}^{k-1} (N_{\alpha, \beta+j\alpha} + N_{\alpha, -\beta-(j+1)\alpha}) e_{\beta+(j+1)\alpha} e_{-\beta-j\alpha} \end{aligned}$$

By the invariance of the bilinear form,

$$\begin{aligned} N_{\alpha, \beta+j\alpha} &= ([e_\alpha, e_{\beta+j\alpha}], e_{-\beta-(j+1)\alpha}) \\ &= -(e_{\beta+j\alpha}, [e_\alpha, e_{-\beta-(j+1)\alpha}]) \\ &= -N_{\alpha, -\beta-(j+1)\alpha}. \end{aligned}$$

Therefore $\sum_{j \in S} [e_\alpha, e_{\beta+j\alpha} e_{-\beta-j\alpha}]$ vanishes. The other statement is proved by replacing α by $-\alpha$ and noticing that $S(-\alpha, \beta) = S(\alpha, \beta)$. \square

Proof of Proposition 2.2: Consider

$$T(\lambda, \mu)T(\nu, \mu) = \sum_{\alpha, \beta} \frac{\langle \alpha, \lambda \rangle \langle \beta, \nu \rangle}{\langle \alpha, \mu \rangle \langle \alpha, \mu \rangle} [e_\alpha e_{-\alpha}, e_\beta e_{-\beta}] \quad (5)$$

Let us show that this expression is a regular function of $\mu \in \mathfrak{h}$. Since for nontrivial λ, μ it converges to zero at infinity, it then vanishes identically.

We compute the residue of (5) at $\langle \alpha, \mu \rangle = 0$:

$$\sum_{\gamma: \gamma \neq \pm\alpha} \frac{\langle \alpha, \lambda \rangle \langle \gamma, \nu \rangle - \langle \gamma, \lambda \rangle \langle \alpha, \nu \rangle}{\langle \gamma, \mu \rangle} [e_\alpha e_{-\alpha}, e_\gamma e_{-\gamma}], \quad (6)$$

a function on the hyperplane $\langle \alpha, \mu \rangle = 0$. The sum over γ of the form $\beta + j\alpha$, $j \in S(\alpha, \beta)$ gives for $\langle \alpha, \mu \rangle = 0$,

$$\sum_{j \in S} \frac{\langle \alpha, \lambda \rangle \langle \beta, \nu \rangle - \langle \beta, \lambda \rangle \langle \alpha, \nu \rangle}{\langle \beta, \mu \rangle} [e_\alpha e_{-\alpha}, e_{\beta+j\alpha} e_{-\beta-j\alpha}] = 0, \quad (7)$$

by the previous Lemma. Since the sum over γ in (6) can be written as a sum of such terms, it vanishes.

3 Hypergeometric solutions

Let \mathfrak{g} be a simple complex Lie algebra. Choose a set $f_1, \dots, f_r, e_1, \dots, e_r$ of Chevalley generators of the Lie algebra \mathfrak{g} associated with simple roots $\alpha_1, \dots, \alpha_r$. Let $\lambda = (m_1, \dots, m_r) \in \mathbb{N}^r$. Let $Q_+ = \sum \mathbb{N}\alpha_i$ be the *positive root lattice* for \mathfrak{g} . Define a map $\alpha : \mathbb{N}^r \rightarrow Q_+$ by $\alpha(\lambda) = \sum m_i \alpha_i$. Let V be a tensor product of highest weight modules V_j of \mathfrak{g} with respective highest weights Λ_j , where $j = 1, \dots, n$. Set $\Lambda = \sum_{j=1}^n \Lambda_j$. Denote V_λ the weight space of V with weight $\Lambda - \alpha(\lambda)$. The hypergeometric solutions of the KZ equations in V_λ , see [SV], have the form $u(z) = \int_{\gamma(z)} \Phi(z, t)^{1/k} \omega(z, t)$. We will describe the explicit construction. The number of integration variables $(t_k)_{k=1}^m$ is $m = \sum m_i$. Let c be the unique non-decreasing function from $\{1, \dots, m\}$ to $\{1, \dots, r\}$ ($i = 1, \dots, r$), such that $\#c^{-1}(\{i\}) = m_i$. Define

$$\Phi(z, t) = \prod_{i < j} (z_i - z_j)^{(\Lambda_i, \Lambda_j)} \prod_{k, j} (t_k - z_j)^{-(\alpha_{c(k)}, \Lambda_j)} \prod_{k < l} (t_k - t_l)^{(\alpha_{c(k)}, \alpha_{c(l)})}.$$

The m -form $\omega(z, t)$ is a closed logarithmic differential form on $\mathbb{C}^n \times \mathbb{C}^m$ with values in V_λ . It has the following combinatorial description. Let $P(\lambda, n)$ be the set of sequences $I = (i_1^1, \dots, i_{s_1}^1; \dots; i_1^n, \dots, i_{s_n}^n)$ of integers in $\{1, \dots, r\}$ with $s_j \geq 0$, $j = 1, \dots, n$ and such that, for all $1 \leq j \leq r$, j appears precisely $|c^{-1}(j)|$ times in I . For $I \in P(\lambda, n)$, and a permutation $\sigma \in \Sigma_m$, set $\sigma_1(l) = \sigma(l)$ and $\sigma_j(l) = \sigma(s_1 + \dots + s_{j-1} + l)$, $j = 2, \dots, n$, $1 \leq l \leq s_j$. Define $\Sigma(I) = \{\sigma \in \Sigma_m \mid c(\sigma_j(l)) = i_j^l \text{ for all } j \text{ and } l\}$.

Fix a highest-weight vector v_j for each representation V_j , $j = 1, \dots, n$. To every $I \in P(\lambda, n)$ we associate a vector $f_I v = f_{i_1^1} \cdots f_{i_{s_1}^1} v_1 \otimes \cdots \otimes f_{i_1^n} \cdots f_{i_{s_n}^n} v_n$ in V_λ , and meromorphic differential m -forms $\omega_{I, \sigma} = \omega_{\sigma_1(1), \dots, \sigma_1(s_1)}(z_1) \wedge \cdots \wedge \omega_{\sigma_n(1), \dots, \sigma_n(s_n)}(z_n)$, labeled by $\sigma \in \Sigma(I)$, where $\omega_{i_1, \dots, i_s}(z) = d \log(t_{i_1} - t_{i_2}) \wedge \cdots \wedge d \log(t_{i_{s-1}} - t_{i_s}) \wedge d \log(t_{i_s} - z)$ is a meromorphic one form on $\mathbb{C} \times \mathbb{C}^s$. Finally

$$\omega(z, t) = \sum_{I \in P(\lambda, n)} \sum_{\sigma \in \Sigma(I)} (-1)^{|\sigma|} \omega_{I, \sigma} f_I v.$$

It obeys the equation $\Phi^{-1} d\Phi \wedge \omega = K \wedge \omega$, where $K = \sum_{i < j} \Omega^{(ij)} \frac{dz_i - dz_j}{z_i - z_j}$. As a consequence, for each horizontal family of twisted cycles $\gamma(z)$ in $\{z\} \times \mathbb{C}^m$, u obeys the KZ equations (3) with $\mu = 0$ (See [SV]). This construction can be modified to give solutions for general μ :

Theorem 3.1 *The integrals*

$$\int_{\gamma(z)} \Phi_\mu^{\frac{1}{k}} \omega, \quad \Phi_\mu = \exp\left(-\sum_{i=1}^m \langle \alpha_{c(i)}, \mu \rangle t_i + \sum_{j=1}^n \langle \Lambda_j, \mu \rangle z_j\right) \Phi, \quad (8)$$

are solutions of the KZ equations (3).

Proof. The proof follows from the identity

$$\Phi_\mu^{-1} d\Phi_\mu \wedge \omega = \sum_{i=1}^n \mu^{(i)} dz_i \wedge \omega + K \wedge \omega.$$

To prove this identity notice that ω is a sum of several terms ω_s , which can be grouped according to how the integration variables t_1, \dots, t_m are distributed among the points z_1, \dots, z_n . If the variable t_k is associated to the point z_i in one term ω_s , then $(dt_k - dz_i) \wedge \omega_s = 0$. Moreover, if the term ω_s obeys $h^{(i)} \omega_s = (\langle \Lambda_i, h \rangle - \sum m'_j(i) \langle \alpha_j, h \rangle) \omega_s$, $h \in \mathfrak{h}$, where Λ_i is the highest weight of the i -th representation, then the number of variables t_k associated to z_i such that $c(k) = j$ is $m'_j(i)$, ($j = 1, \dots, r$). Thus we have

$$\Phi_\mu^{-1} d\Phi_\mu \wedge \omega = \sum_{i=1}^n (\langle \Lambda_i, \mu \rangle - \sum_j m'_j(i) \langle \alpha_j, \mu \rangle) dz_i \wedge \omega + K \wedge \omega.$$

The proof is complete. □

Theorem 3.2 *The hypergeometric integrals of Theorem 3.1 obey the Dynamical differential equations (4).*

The proof of the Theorem is given in Section 7 . □

An application of the above two theorems is a determinant formula.

Corollary 3.3 *Fix a basis v_1, \dots, v_d of a weight space V_λ . Suppose that $u_i(\mu, z) = \sum_{j=1}^d u_{i,j} v_j$, $i = 1, \dots, d$ is a basis of the space of solutions in a neighbourhood of a generic point $(\mu, z) \in \mathfrak{h} \times \mathbb{C}^n$. Let $\delta_\alpha = \text{tr}_{V_\lambda}(e_{-\alpha} e_\alpha)$, ($\alpha \in \Delta$, $\alpha > 0$), $\epsilon_{ij} = \text{tr}_{V_\lambda}(\Omega_{ij})$. Then there is a constant $C = C(V_1, \dots, V_n, \lambda, \kappa) \neq 0$ such that*

$$\det(u_{ij}) = C \exp\left(\sum_{i=1}^n \frac{z_i}{\kappa} \text{tr}_{V_\lambda}(\mu^{(i)})\right) \prod_{\alpha > 0} \langle \alpha, \mu \rangle^{\frac{\delta_\alpha}{\kappa}} \prod_{i < j} (z_i - z_j)^{\frac{\epsilon_{ij}}{\kappa}}.$$

4 Free Lie algebras. Dynamical differential equations

4.1 The definition of KZ and dynamical differential equations.

Following [SV] let us fix the following data:

1. A finite dimensional complex vector space \mathfrak{h} ;
2. A non-degenerate symmetric bilinear form $(,)$ on \mathfrak{h} ;
3. Linearly independent covectors (“simple roots”) $\alpha_1, \dots, \alpha_r \in \mathfrak{h}^*$.

We denote by $b : \mathfrak{h} \rightarrow \mathfrak{h}^*$ the isomorphism induced by $(,)$, and we transfer the form $(,)$ to \mathfrak{h}^* via b . Set $b_{ij} = (\alpha_i, \alpha_j)$, $h_i = b^{-1}(\alpha_i) \in \mathfrak{h}$. Denote by \mathfrak{g} the Lie algebra generated by e_i, f_i for $i = 1, \dots, r$ and \mathfrak{h} subject to the relations:

$$[h, e_i] = \langle \alpha_i, h \rangle e_i, \quad [h, f_i] = -\langle \alpha_i, h \rangle f_i, \quad [e_i, f_j] = \delta_{ij} h_i, \quad [h, h'] = 0,$$

for all $i, j = 1, \dots, r$ and $h, h' \in \mathfrak{h}$. Thus we have constructed a Kac-Moody Lie algebra without Serre’s relations. We denote by \mathfrak{n}_- (resp. by \mathfrak{n}_+) the subalgebra of \mathfrak{g} generated by f_i (resp. e_i) for $i = 1, \dots, r$. We have $\mathfrak{g} = \mathfrak{n}_- \oplus \mathfrak{h} \oplus \mathfrak{n}_+$. Set $\mathfrak{b}_\pm = \mathfrak{n}_\pm \oplus \mathfrak{h}$. These are subalgebras of \mathfrak{g} .

Let $\Lambda \in \mathfrak{h}^*$. Denote by $M(\Lambda)$ the Verma module over \mathfrak{g} generated by a vector v subject to the relations $\mathfrak{n}_+ v = 0$ and $h v = \langle \Lambda, h \rangle v$ for all $h \in \mathfrak{h}$. Let us fix weights $\Lambda_1, \dots, \Lambda_n \in \mathfrak{h}^*$, and let $M = M(\Lambda_1) \otimes \dots \otimes M(\Lambda_n)$.

For $\lambda = (m_1, \dots, m_r) \in \mathbb{N}^r$, set

$$\begin{aligned} (\mathfrak{n}_\pm)_\lambda &= \left\{ x \in \mathfrak{n}_\pm \mid [h, x] = \langle \pm \sum m_i \alpha_i, h \rangle x, \quad \text{for all } h \in \mathfrak{h} \right\}, \\ M(\Lambda)_\lambda &= \left\{ x \in M(\Lambda) \mid h x = \langle \Lambda - \sum m_i \alpha_i, h \rangle x \quad \text{for all } h \in \mathfrak{h} \right\}, \\ M_\lambda &= \left\{ x \in M \mid h x = \langle \sum \Lambda_j - \sum m_i \alpha_i, h \rangle x \quad \text{for all } h \in \mathfrak{h} \right\}. \end{aligned}$$

We have $\mathfrak{n}_\pm = \bigoplus_\lambda (\mathfrak{n}_\pm)_\lambda$, $\mathfrak{b}_\pm = \mathfrak{h} \oplus (\bigoplus_\lambda (\mathfrak{n}_\pm)_\lambda)$, $M(\Lambda) = \bigoplus_\lambda M(\Lambda)_\lambda$, $M = \bigoplus_\lambda M_\lambda$.

Let $\tau : \mathfrak{g} \rightarrow \mathfrak{g}$ be the Lie algebra automorphism such that $\tau(e_i) = -f_i$, $\tau(f_i) = -e_i$, $\tau(h) = -h$, for $h \in \mathfrak{h}$. Set $\mathfrak{n}_\pm^* = \bigoplus_\lambda (\mathfrak{n}_\pm)_\lambda^*$. Set $M(\Lambda)^* = \bigoplus_\lambda M(\Lambda)_\lambda^*$. Define a structure of a

\mathfrak{g} -module on $M(\Lambda)^*$ by the rule

$$\langle g\phi, x \rangle = \langle \phi, -\tau(g)x \rangle \text{ for } \phi \in M(\Lambda)^*, \quad g \in \mathfrak{g}, \quad x \in M(\Lambda). \quad (9)$$

There is a unique bilinear form $K(\cdot, \cdot)$ on \mathfrak{g} such that: K coincides with (\cdot, \cdot) on \mathfrak{h} ; K is zero on \mathfrak{n}_+ and \mathfrak{n}_- ; \mathfrak{h} and $\mathfrak{n}_- \oplus \mathfrak{n}_+$ are orthogonal; $K(f_i, e_j) = K(e_j, f_i) = \delta_{ij}$ for $i, j = 1, \dots, r$; K is \mathfrak{g} -invariant, that is $K([x, y], z) = K(x, [y, z])$ for all $x, y, z \in \mathfrak{g}$.

A bilinear form S on \mathfrak{g} is defined by the rule $S(x, y) = -K(\tau(x), y)$. The form S is symmetric, τ -invariant, and $S([x, y], z) = S(x, [\tau(y), z])$. The subspaces \mathfrak{n}_+ , \mathfrak{h} , \mathfrak{n}_- are pairwise orthogonal with respect to S .

For a Verma module $M(\Lambda)$ with highest weight Λ and generating vector v there is a unique bilinear form S on $M(\Lambda)$ such that: $S(v, v) = 1$; $S(e_i x, y) = S(x, f_i y)$; $S(f_i x, y) = S(x, e_i y)$, for all $x, y \in M(\Lambda)$ and $i = 1, \dots, r$. S is symmetric. The subspaces $M(\Lambda)_\lambda$ are pairwise orthogonal with respect to S . The form S induces a homomorphism of \mathfrak{g} -modules $S : M(\Lambda) \rightarrow M(\Lambda)^*$. The module $M(\Lambda)/\ker S$ is the irreducible \mathfrak{g} -module with highest weight Λ . More generally, on the space $\wedge^p \mathfrak{n}_- \otimes M(\Lambda_1) \otimes \dots \otimes M(\Lambda_n)$ for $\Lambda_j \in \mathfrak{h}^*$, a bilinear form S is defined by the rule

$$S(g_1 \wedge \dots \wedge g_p \otimes x_1 \otimes \dots \otimes x_n, g'_1 \wedge \dots \wedge g'_p \otimes x'_1 \otimes \dots \otimes x'_n) = \det S(g_i, g'_j) \cdot \prod_{i=1}^n S(x_i, x'_i).$$

This form induces a map

$$MS : \wedge^p \mathfrak{n}_- \otimes M(\Lambda_1) \otimes \dots \otimes M(\Lambda_n) \rightarrow (\wedge^p \mathfrak{n}_- \otimes M(\Lambda_1) \otimes \dots \otimes M(\Lambda_n))^* \quad (10)$$

S is called the contravariant form. The linear map (10) depends analytically on (b_{ij}) and $\Lambda_1, \dots, \Lambda_n \in \mathfrak{h}^*$ for a fixed r . It is non-degenerate for general values of the parameters, see Theorem 3.7 in [SV].

A Lie bialgebra structure on \mathfrak{b}_\pm [SV]. A Lie bialgebra is a vector space \mathfrak{g} with a Lie algebra structure and a Lie coalgebra structure, such that the cocommutator map $\nu : \mathfrak{g} \rightarrow \mathfrak{g} \wedge \mathfrak{g}$ is a one-cocycle: $x\nu(y) - y\nu(x) = \nu([x, y])$, for all $x, y \in \mathfrak{g}$. Here the action of \mathfrak{g} on $\mathfrak{g} \wedge \mathfrak{g}$ is the adjoint one: $a(b \wedge c) = [a, b] \wedge c + b \wedge [a, c]$. The dual map to the cocommutator map, $\nu^* : (\mathfrak{g} \wedge \mathfrak{g})^* \rightarrow \mathfrak{g}^*$, defines a Lie algebra structure on \mathfrak{g}^* .

Let \mathfrak{g} be a Lie bialgebra. The double of \mathfrak{g} is the Lie algebra equal to $\mathfrak{g} \oplus \mathfrak{g}^*$ as a vector space with the bracket on \mathfrak{g} and \mathfrak{g}^* defined by the Lie algebra structure on \mathfrak{g} and \mathfrak{g}^* , and for $x \in \mathfrak{g}$ and $l \in \mathfrak{g}^*$, $[l, x] = \bar{l} + \bar{x}$, where $\bar{x} \in \mathfrak{g}$ and $\bar{l} \in \mathfrak{g}^*$ are defined by the rules $\bar{l}(y) = l([x, y])$, $m(\bar{x}) = [m, l](x)$, for all $y \in \mathfrak{g}$, $m \in \mathfrak{g}^*$. The double is denoted by $D(\mathfrak{g})$.

Let \mathfrak{g} be the Kac-Moody algebra we defined at the beginning of the section. \mathfrak{g} is a Lie bialgebra with respect to the following cobracket. There exists a unique map $\nu : \mathfrak{g} \rightarrow \mathfrak{g} \wedge \mathfrak{g}$ such that $x\nu(y) - y\nu(x) = \nu([x, y])$, and $\nu(h) = 0$, and $\nu(f_i) = \frac{1}{2}f_i \wedge h_i$, and $\nu(e_i) = \frac{1}{2}e_i \wedge h_i$. In the previous four equalities, $h \in \mathfrak{h}$, $i = 1, \dots, r$, the action of \mathfrak{g} on $\mathfrak{g} \wedge \mathfrak{g}$ is the adjoint one, see [D] Example 3.2 and [SV]. \mathfrak{b}_- and \mathfrak{b}_+ are subbialgebras. The map ν has the property $\tau\nu + \nu\tau = 0$. Thus, if $\rho : \mathfrak{b}_-^* \rightarrow \text{End}(V)$ is a representation of the Lie algebra

$(\nu|_{\mathfrak{b}_-})^* : \Lambda^2 \mathfrak{b}_-^* \rightarrow \mathfrak{b}_-^*$, then $-\rho \circ \tau : \mathfrak{b}_+^* \rightarrow \text{End}(V)$ is a representation of the Lie algebra $(\nu|_{\mathfrak{b}_+})^* : \Lambda^2 \mathfrak{b}_+^* \rightarrow \mathfrak{b}_+^*$.

Note that the coalgebra map ν defines a Lie algebra structure on \mathfrak{b}_\pm^* .

Comultiplication. Let $M = M(\Lambda_1) \otimes \cdots \otimes M(\Lambda_n)$. Let $v = v_1 \otimes \cdots \otimes v_n \in M$ be the product of the generating vectors. Set $\Lambda = \sum \Lambda_j$. Let \mathfrak{b}_- act on $\mathfrak{b}_- \otimes M$ by the rule $a(b \otimes m) = [a, b] \otimes m + b \otimes am$. For $1 \leq i \leq n$, and $a, b \in \mathfrak{g}$, $m = x_1 \otimes \cdots \otimes x_n \in M$, set $a^{(i)}m = x_1 \otimes \cdots \otimes x_{i-1} \otimes ax_i \otimes x_{i+1} \otimes \cdots \otimes x_n$ and $a^{(i)}(b \otimes m) = [a, b] \otimes m + b \otimes a^{(i)}m$.

There is a unique linear map $\nu_M : M \rightarrow \mathfrak{b}_- \otimes M$ such that

$$\begin{aligned} \nu_M(h \cdot x) &= h \cdot \nu_M(x) \text{ for any } h \in \mathfrak{h} \text{ and } x \in M; \\ \nu_M(x) &= \frac{1}{2}(b^{-1}(\Lambda - \alpha(\lambda))) \otimes x + \nu_{M-}(x) \text{ for } x \in M_\lambda. \end{aligned}$$

Recall that $b^{-1}(\alpha(\lambda)) = \sum m_i b^{-1}(\alpha_i) = \sum m_i h_i$, and $b^{-1} : \mathfrak{h}^* \rightarrow \mathfrak{h}$ is defined at the beginning of Sect. 4.1. The map $\nu_{M-} : M \rightarrow \mathfrak{n}_- \otimes M$ is defined via an inductive definition. $\nu_{M-}(v) = 0$, $\nu_{M-}(x) = \sum_{k=1}^n \nu_{M-}^{(k)}(x)$ where

$$\nu_{M-}^{(k)}(f_i^{(j)}x) = f_i^{(j)} \nu_{M-}^{(k)}(x) \text{ for } k \neq j, \text{ and } \nu_{M-}^{(k)}(f_i^{(k)}x) = f_i \otimes h_i^{(k)}x + f_i^{(k)} \nu_{M-}^{(k)}(x).$$

In all formulae we have $1 \leq i \leq r$, and $1 \leq j, k \leq n$.

Remark. The corresponding definition of $\nu_{M-}(x)$ in [SV] should be corrected as above. Note that the two definitions coincide if we have one tensor factor, i.e. $n = 1$. We have the following Lemma (cf. Lemma 6.15.2 in [SV]).

Lemma 4.1 *For any $x, y \in M$, $a \in \mathfrak{b}_-$,*

$$S(\nu_M(x), a \otimes y) = \begin{cases} \frac{1}{2}S(x, ay) & \text{if } a \in \mathfrak{h}; \\ S(x, ay) & \text{if } a \in \mathfrak{n}_-. \end{cases} \quad (11)$$

Here S is defined on $\mathfrak{b}_- \otimes M$ by the rule $S(a \otimes x, b \otimes y) = S(a, b)S(x, y)$, cf. (10).

The proof of Lemma 4.1 is given in Section 4.5 □

Note that the above Lemma renders the following diagram commutative.

$$\begin{array}{ccc} \mathfrak{n}_- \otimes M & \xrightarrow{\text{standard}} & M \\ \downarrow S & & \downarrow S \\ \mathfrak{n}_-^* \otimes M^* & \xrightarrow{\nu_{M-}^*} & M^* \end{array}$$

S is an isomorphism for general values of parameters (b_{ij}) , $(\Lambda_k)_{k=1}^n$, see [SV] sec. (3.7) and (6.6). Hence, $\nu_M^* : \mathfrak{b}_-^* \otimes M^* \rightarrow M^*$ is a \mathfrak{b}_-^* -module structure with respect to the Lie algebra structure $\nu^* : \Lambda^2 \mathfrak{b}_-^* \rightarrow \mathfrak{b}_-^*$ for any values of the above parameters.

Corollary 4.2 *If $y \in \ker S : \mathfrak{n}_- \rightarrow \mathfrak{n}_-$ then $yM \subset \ker S : M \rightarrow M$.* □

Actions of the doubles of \mathfrak{b}_\pm on M and M^* respectively.

Consider the standard action of \mathfrak{b}_- on M^* , i.e. $\forall a \in \mathfrak{b}_- \forall \phi \in M^*, \langle a \cdot \phi, \cdot \rangle = \langle \phi, -a \cdot \rangle$ where the action on the right hand side is the standard action of \mathfrak{g} on M . This map together with $\nu_{M^-}^*$ defines an action of $\mathfrak{b}_- \oplus \mathfrak{b}_-^*$ on M^* . Lemma 6.17.1 [SV] asserts that M^* is a $D(\mathfrak{b}_-)$ -module under this action, where $D(\mathfrak{b}_-)$ is the double of \mathfrak{b}_- .

For any $a \in \mathfrak{b}_-^*$, the action of ν_M^* defines a map $\nu_M^*(a, \cdot) : M^* \rightarrow M^*$. Set $\rho(a) = -(\nu_M^*(a, \cdot))^* : M \rightarrow M$. The map $\rho : \mathfrak{b}_-^* \rightarrow \text{End}(M)$ gives an action of \mathfrak{b}_-^* on M . An action of \mathfrak{b}_+^* on M is defined by $\omega = -\rho \circ \tau : \mathfrak{b}_+^* \rightarrow \text{End}(M)$. The rule $a \otimes x \rightarrow \tau(a)x$, for $a \in \mathfrak{b}_+, x \in M$, defines an action of \mathfrak{b}_+ on M . This action and the map ω define an action of $\mathfrak{b}_+ \oplus \mathfrak{b}_+^*$ on M . Lemma 11.3.28 [V] implies that M is a $D(\mathfrak{b}_+)$ module.

Note that for any Kac-Moody Lie algebra \mathfrak{g} without Serre's relations, and a \mathfrak{g} -module V we can define a \mathfrak{g} -module structure on V^* by the rule $\langle g\phi, \cdot \rangle = \langle \phi, -g \cdot \rangle$ for all $g \in \mathfrak{g}$ and $\phi \in M^*$. With respect to this module structure, M becomes a $D(\mathfrak{b}_-)$ -module and M^* becomes a $D(\mathfrak{b}_+)$ -module.

KZ equations and dynamical differential equations in M and M^* . For a vector space V denote by $\Omega(V) \in V \otimes V^*$ the canonical element. For $\lambda \in \mathbb{N}^r$, set $\Omega_{\lambda, \pm}^- := \Omega((\mathfrak{n}_\pm)_\lambda) \in (\mathfrak{n}_\pm)_\lambda \otimes (\mathfrak{n}_\pm)_\lambda^*$, set $\Omega_{\lambda, \pm}^+ := \Omega((\mathfrak{n}_\pm)_\lambda^*) \in (\mathfrak{n}_\pm)_\lambda^* \otimes (\mathfrak{n}_\pm)_\lambda$, $\Omega^0 := (\Omega(\mathfrak{h}) + \Omega(\mathfrak{h}^*)) \in \mathfrak{h} \otimes \mathfrak{h}^* + \mathfrak{h}^* \otimes \mathfrak{h}$. Set

$$\Omega_\pm = \sum_\lambda \Omega_{\lambda, \pm}^- + \Omega^0 + \sum_\lambda \Omega_{\lambda, \pm}^+ \in D(\mathfrak{b}_\pm) \otimes D(\mathfrak{b}_\pm).$$

Let $\Omega_{+,ij}$ be the operator on M (or M^*) acting as Ω_+ on $M(\Lambda_i) \otimes M(\Lambda_j)$, ($M(\Lambda_i)^* \otimes M(\Lambda_j)^*$ respectively) and as the identity on the other factors. The action of $D(\mathfrak{b}_\pm)$ on $M(\Lambda_j)$ is the action described in the previous chapter when $n = 1$. The KZ equations with additional parameter $\mu \in \mathfrak{h}$, for a function $u(\mu, z)$ on n variables $z = (z_1, \dots, z_n)$ taking values in M (or M^*) are

$$\kappa \frac{\partial u}{\partial z_i} = \mu^{(i)} u + \sum_{i \neq j} \frac{\Omega_{+,ij}}{z_i - z_j} u, \quad i = 1, \dots, n, \quad \kappa \in \mathbb{C}. \quad (12)$$

Let α be a positive root for \mathfrak{g} and $(y_i^{(\alpha)})$ a basis of $(\mathfrak{n}_+)_\alpha$. Set $x_i^{(\alpha)} = \tau(y_i^{(\alpha)})$. Then $((y_i^{(\alpha)})^*)$, $(x_i^{(\alpha)})$ and $((x_i^{(\alpha)})^*)$ are bases of $(\mathfrak{n}_+^*)_\alpha$, $(\mathfrak{n}_-)_\alpha$, $(\mathfrak{n}_-^*)_\alpha$, respectively. Define operators $\Delta_{\pm, \alpha}$ on M (or M^*) via the formulae $\Delta_{+, \alpha} = \sum_i y_i^{(\alpha)} (y_i^{(\alpha)})^*$, $\Delta_{-, \alpha} = \sum_i (x_i^{(\alpha)})^* x_i^{(\alpha)}$. The dynamical differential equations for the function $u(\mu, z)$ with values in M (M^* respectively) are

$$\kappa \partial_{\mu'} u = \sum_{i=1}^n z_i (\mu')^{(i)} u + \sum_{\alpha > 0} \frac{\langle \alpha, \mu' \rangle}{\langle \alpha, \mu \rangle} \Delta_{+, \alpha} u, \quad \mu' \in \mathfrak{h}, \quad \kappa \in \mathbb{C}. \quad (13)$$

4.2 Properties of the operators $\Delta_{+, \alpha}$.

The properties of the operator $\Omega_{+,ij}$ are thoroughly described in [V]. Now we are going to study the operators $\Delta_{+, \alpha}$.

Lemma 4.3 *The following diagram is commutative:*

$$\begin{array}{ccc} M & \xrightarrow{\Delta_{+,\alpha}} & M \\ \downarrow S & & \downarrow S \\ M^* & \xrightarrow{-\Delta_{-,\alpha}} & M^* \end{array}$$

In particular, the operators $\Delta_{+,\alpha}$ preserve the kernel of the map $S : M \rightarrow M^*$.

Proof. We fix α throughout this proof and will drop it from the notation of the bases. Fix a basis (u_k) of $\ker S : (\mathbf{n}_+)_{\alpha} \rightarrow (\mathbf{n}_+^*)_{\alpha}$. Complete it to a basis of $(\mathbf{n}_+)_{\alpha}$ by vectors (v_l) . Let $(u_k^*), (v_l^*)$ be the dual basis of $(\mathbf{n}_+^*)_{\alpha}$. Moreover,

$$(v_l^*) = \sum_p (A^{-1})_{lp} S(v_p, \cdot), \quad \text{and} \quad (\tau(v_l)^*) = \sum_p (A^{-1})_{lp} S(\tau(v_p), \cdot), \quad (14)$$

where $A = (a_{lp})$ is a nondegenerate matrix with entries $a_{lp} = S(v_l, v_p)$.

For $y \in (\mathbf{n}_+)_{\alpha}$, consider the map $y : M \rightarrow M$ via the action of $D(\mathbf{b}_+)$. Let $y \cdot p$ denote the $D(\mathbf{b}_+)$ action, and yp denote the standard action of \mathbf{b}_+ . Then we have

$$S(y \cdot p, q) = S(\tau(y)p, q) = S(p, -yq) \quad \text{for any } p, q \in M. \quad (15)$$

Consider the map $(v_l)^* : M \rightarrow M$ via the action of $D(\mathbf{b}_+)$. For any $p, q \in M$ we have

$$\begin{aligned} S((v_l)^*p, q) &= \langle (v_l)^*p, S(q, \cdot) \rangle = \langle (\nu_M^*(\tau(v_l)^*, \cdot))^*p, S(q, \cdot) \rangle \\ &= \langle p, \nu_M^*(\tau(v_l)^*, S(q, \cdot)) \rangle = \langle p, \nu_M^*(\sum_j (A^{-1})_{lj} S(\tau(v_j), \cdot), S(q, \cdot)) \rangle \\ &= \sum_j (A^{-1})_{lj} \langle p, S(\tau(v_j)q, \cdot) \rangle = \sum_j (A^{-1})_{lj} S(p, \tau(v_j)q). \end{aligned} \quad (16)$$

The first three equalities come from the definition of the action of $D(\mathbf{b}_+)$ on M , the last two from formula (14). We combine (15), (16), and Corollary 4.2 to obtain:

$$\begin{aligned} S(\Delta_{+,\alpha}p, q) &= S((\sum_k u_k(u_k^*) + \sum_l v_l(v_l^*))p, q) = S(\sum_l v_l(v_l^*)p, q) \\ &= S((v_l^*)p, (-v_l)q) = -\sum_{j,l} (A^{-1})_{lj} S(p, \tau(v_j)v_lq). \end{aligned} \quad (17)$$

Now we trace the arrows in the alternative direction. For $x \in (\mathbf{n}_-)_{\alpha}$, consider the map $x : M^* \rightarrow M^*$ via the action of $D(\mathbf{b}_-)$. Denote this action by \cdot' . Let xq denote the standard action of \mathbf{b}_- on M . We have

$$\langle x \cdot S(p, \cdot), q \rangle = \langle S(p, -\tau(\tau(x)) \cdot), q \rangle = S(p, -xq) \quad \text{for all } x \in \mathbf{b}_-; p, q \in M. \quad (18)$$

Consider the map $(\tau(v_l))^* : M^* \rightarrow M^*$ via the action of $D(\mathbf{b}_-)$.

$$\begin{aligned} \langle (\tau(v_l))^* S(p, \cdot), q \rangle &= \langle (\sum_j (A^{-1})_{lj} S(\tau(v_j), \cdot)) S(p, \cdot), q \rangle \\ &= \sum_j (A^{-1})_{lj} \langle S(\tau(v_j)p, \cdot), q \rangle = \sum_j (A^{-1})_{lj} S(\tau(v_j)p, q). \end{aligned} \quad (19)$$

Finally combine (18), (19), and Corollary 4.2 with $u_k \in \ker S$ to get

$$\begin{aligned} \langle -\Delta_{-, \alpha} S(p, \cdot), q \rangle &= \langle - \left(\sum_k \tau(u_k)^* \tau(u_k) + \sum_l \tau(v_l)^* \tau(v_l) \right) S(p, \cdot), q \rangle \\ &= - \langle \sum_k \tau(u_k)^* S(p, -\tau(u_k) \cdot), q \rangle - \langle \sum_l \tau(v_l)^* S(p, -\tau(v_l) \cdot), q \rangle \\ &= \langle - \sum_k \tau(u_k)^* S(-\tau(-\tau(u_k))p, \cdot), q \rangle + \langle - \sum_l \tau(v_l)^* S(-\tau(-\tau(v_l))p, \cdot), q \rangle \\ &= 0 - \langle \sum_{j,l} (A^{-1})_{lj} S(\tau(v_j)(v_l)p, \cdot), q \rangle = - \sum_{j,l} (A^{-1})_{lj} S(\tau(v_j)(v_l)p, q) \\ &= - \sum_{j,l} (A^{-1})_{lj} S(p, \tau(v_l)v_j q). \end{aligned} \quad (20)$$

Since the matrix A is symmetric (17) and (20) prove that the diagram is commutative. \square

As a corollary of the Lemma we have that $\Delta_{+, \alpha}$ naturally acts on $L = M / \ker(S : M \rightarrow M^*)$. We describe this action. Consider the Kac-Moody algebra $\bar{\mathfrak{g}} = \mathfrak{g} / \ker(S : \mathfrak{g} \rightarrow \mathfrak{g}^*)$. Let $x \mapsto \bar{x}$ denote the canonical projections $M \rightarrow L$, $\mathfrak{g} \rightarrow \bar{\mathfrak{g}}$. $\ker S$ is an ideal and the form S induces a non-degenerate Killing form on $\bar{\mathfrak{g}}$ via the formula $K(x, y) = -S(\tau(x), y)$, see [V]. K induces a non-degenerate pairing between root spaces $\bar{\mathfrak{g}}_\alpha$ and $\bar{\mathfrak{g}}_{-\alpha}$. Let $(e_l^{(\alpha)})$ be a basis of $\bar{\mathfrak{g}}_\alpha$, and let $(f_l^{(\alpha)})$ be the dual basis of $\bar{\mathfrak{g}}_{-\alpha}$ with respect to K . Let $\bar{\Delta}_\alpha = \sum_l f_l^{(\alpha)} e_l^{(\alpha)}$.

Corollary 4.4 *The following diagram is commutative*

$$\begin{array}{ccc} M & \xrightarrow[\substack{D(\mathbf{b}_+)-\text{action}}]{\Delta_{+, \alpha}} & M \\ \downarrow & & \downarrow \\ L & \xrightarrow[\substack{\text{standard}}]{\bar{\Delta}_\alpha} & L. \end{array}$$

Proof. $L(\Lambda) \cong \text{Im}\{S : M \rightarrow M^*\}$ via $\bar{x} \mapsto S(x, \cdot)$. We keep the notation from the Lemma above. Set $w_l = - \sum_j (A^{-1})_{lj} \tau(v_j)$. From the computation in the Lemma we have

$$\begin{aligned} \overline{\Delta_{+, \alpha} p} &= S(\Delta_{+, \alpha} p, \cdot) = S(\sum_l (- \sum_j (A^{-1})_{lj} \tau(v_j)) v_l p, \cdot) \\ &= S(\sum_l w_l v_l p, \cdot) = \overline{\sum_l w_l v_l p} = \sum_l \bar{w}_l \bar{v}_l \bar{p} \end{aligned}$$

Finally notice that the set (\bar{v}_i) forms a basis of $\bar{\mathfrak{g}}_\alpha$, and the set (\bar{w}_i) forms the dual basis of $\bar{\mathfrak{g}}_{-\alpha}$ with respect to K . \square

Corollary 4.5 *Fix $\lambda \in \mathbb{N}^r$. Let $m \in M_\lambda$, and let (m_j) be a basis of M_λ , and let (m_j^*) be the dual basis of M_λ^* . Then the following decomposition holds*

$$\Delta_{+,\alpha}m = \sum_j \langle -\Delta_{-,\alpha}m_j^*, m \rangle m_j.$$

Proof. Let $y \in \mathfrak{b}_+$ and $x = \tau(y) \in \mathfrak{b}_-$. Let $p \in M$, and $\phi \in M^*$. As in the proof of Lemma 4.3 $\langle y^*p, \phi \rangle = \langle p, x^*\phi \rangle$, where $D(\mathfrak{b}_+)$ acts on M and $D(\mathfrak{b}_-)$ acts on M^* . Moreover $\langle y \cdot p, \phi \rangle = \langle p, -x \cdot \phi \rangle$, where $D(\mathfrak{b}_+)$ acts on M and $D(\mathfrak{b}_-)$ acts on M^* . Finally noting that $\Delta_{+,\alpha} = \sum_i y_i^{(\alpha)} (y_i^{(\alpha)})^*$ and $\Delta_{-,\alpha} = \sum_i \tau(y_i^{(\alpha)})^* \tau(y_i^{(\alpha)})$ we have

$$\Delta_{+,\alpha}m = \sum_j \langle m_j^*, \Delta_{+,\alpha}m \rangle m_j = \sum_j \langle -\Delta_{-,\alpha}m_j^*, m \rangle m_j. \quad \square$$

4.3 An integral form of the dynamical differential equations

Our aim now is to rewrite the Dynamical equations in a form related to the hypergeometric solutions. Fix $\lambda = (m_1, \dots, m_r) \in \mathbb{N}^r$. Let M be a tensor product of highest weight modules of the Kac-Moody Lie algebra without Serre's relations \mathfrak{g} . Set $\Lambda = \sum \Lambda_j$, the sum of the respective highest weights. Consider the weight space M_λ of M with weight $\Lambda - \alpha(\lambda)$, where $\alpha(\lambda) = \sum m_i \alpha_i$. Fix a highest-weight vector v_j for each module $M(\Lambda_j)$, $j = 1, \dots, n$. To every $I \in P(\lambda, n)$ we associate a vector $f_I v = f_{i_1} \cdots f_{i_{s_1}} v_1 \otimes \cdots \otimes f_{i_n} \cdots f_{i_{s_n}} v_n$ in M_λ , cf. Section 3. Note that the vectors $(f_I v)_{I \in P(\lambda, n)}$ form a basis of M_λ .

\mathfrak{n}_- acts on M_λ^* via the $D(\mathfrak{b}_-)$ action. Therefore $U(\mathfrak{n}_-)$ acts on M_λ^* . Explicitly, $x \cdot \phi(\cdot) = \phi(-x \cdot \cdot)$ for $x \in \mathfrak{n}_-$, $\phi \in M^*$ (cf. Section 4.1), where the action on the left hand side is the $D(\mathfrak{b}_-)$ one and the action on the right hand side is the standard one. Let V be a vector space freely generated by f_1, \dots, f_r . Therefore we have an inclusion of tensor algebras $T(V) \subset T(\mathfrak{n}_-)$. Moreover $T(V)$ is an associative enveloping algebra of the Lie algebra \mathfrak{n}_- . Since $T(V)$ is a free associative algebra, $T(V)$ is isomorphic to the universal enveloping algebra $U(\mathfrak{n}_-)$. From now on we will refer to the monomial basis of $T(V)$ as to the monomial basis of $U(\mathfrak{n}_-)$, and to the dual of the monomial basis of $T(V)$ as to the monomial basis of $U(\mathfrak{n}_-)^*$. Rewrite a commutator $x \in \mathfrak{n}_-$ as an element of $U(\mathfrak{n}_-)$ in the form $x = \sum a_j x_j$ where $a_j \in \mathbb{Z}$, and x_j 's are elements of the monomial basis of $U(\mathfrak{n}_-)$. Thus

$$\langle x \cdot \phi, \cdot \rangle = \langle \phi, - \sum a_j x_j \cdot \rangle \quad (21)$$

Denote $i : \mathfrak{n}_- \rightarrow U(\mathfrak{n}_-)$ the inclusion monomorphism. Let $\sigma_j \in \{1, \dots, r\}$ for $j = 1, \dots, k$. Let the positive root of \mathfrak{g} , $\sum_{j=1}^n \alpha_{\sigma_j}$, correspond to the r -tuple $\lambda' \in \mathbb{N}^r$, i.e. $\alpha(\lambda') = \sum_{j=1}^n \alpha_{\sigma_j}$. Define an element $\Delta_{\sigma_1, \dots, \sigma_k}$ of $(\mathfrak{n}_-)^*_{\lambda'}$ via the rule

$$\langle \Delta_{\sigma_1, \dots, \sigma_k}, x \rangle = \langle (f_{\sigma_1} \cdots f_{\sigma_k})^*, i(x) \rangle \text{ where } x \in \mathfrak{n}_-, \text{ and } (f_{\sigma_1} \cdots f_{\sigma_k})^* \in U(\mathfrak{n}_-)^*. \quad (22)$$

Thus $\langle \Delta_{\sigma_1, \dots, \sigma_k}, x \rangle$ is the coefficient of $f_{\sigma_1} \cdots f_{\sigma_k}$ in the decomposition of $i(x)$ into a sum of monomials .

Lemma 4.6 *Let $\alpha = \sum_{k=1}^r m'_k \alpha_k$ be a positive root of \mathfrak{g} . Let $I \in P(\lambda, n)$. Set $\lambda' = (m'_1, \dots, m'_r) \in \mathbb{N}^r$. Then we have*

$$-\Delta_{-, \alpha}(f_I v)^* = \left(\sum_{(i_1, \dots, i_{m'}) \in P(\lambda', 1)} \Delta_{i_1, \dots, i_{m'}}(e_{i_{m'}} \cdots e_{i_1}) \right) (f_I v)^*, \quad (23)$$

where $\Delta_{i_1, \dots, i_{m'}}$ acts according to the $D(\mathbf{n}_-)$ action on M_λ^* , the product of e 's acts on one tensor factor at a time $(e_{i_{m'}} \cdots e_{i_1}) = \sum_{j=1}^n e_{i_{m'}}^{(j)} \cdots e_{i_1}^{(j)}$, and e_j acts via the standard action (9) on each tensor factor.

Proof. Note that $\alpha = \alpha(\lambda')$. Let $x \in (\mathbf{n}_-)_{\lambda'}$. If there exists i such that $m'_i > m_i$, then formula (21) implies $x \cdot (f_I v)^* = 0$ for every $I \in P(\lambda, n)$ because each monomial x_j in the $U(\mathbf{n}_-)$ expansion of x has more f_i 's than $f_I v$.

Now let $m'_i \leq m_i$ for any $1 \leq i \leq r$. Set $m' = \sum m'_i$. First consider the case $n = 1$, $I = (i_1, \dots, i_{m'}) \in P(\lambda, 1)$. Let (x_j^*) be a basis of $(\mathbf{n}_-)_{\lambda'}$ such that $x_1^* = \Delta_{i_1, \dots, i_{m'}}$, and let (x_j) be the dual basis of $(\mathbf{n}_-)_{\lambda'}$. Formula (22) implies that the coefficient of the monomial $f_{i_1} \cdots f_{i_{m'}}$ in the $U(\mathbf{n}_-)$ expansion of x_1 is 1, and the coefficient of $f_{i_1} \cdots f_{i_{m'}}$ in the $U(\mathbf{n}_-)$ expansion of x_j is 0 for $j > 1$. Now use (21) to obtain

$$\begin{aligned} -x_1 \cdot (f_I v)^* &= (f_{i_{m'+1}} \cdots f_{i_m} v)^* = e_{i_{m'}} \cdots e_{i_1} (f_I v)^* \\ -x_j \cdot (f_I v)^* &= 0 \quad \text{for } j \geq 1. \end{aligned} \quad (24)$$

For any element $I' = (i'_1, \dots, i'_{m'}) \in P(\lambda', 1)$, such that $I' \neq (i_1, \dots, i_{m'})$ we have $e_{i'_{m'}} \cdots e_{i'_1} (f_I v)^* = 0$. The proof for $n = 1$ is finished.

Let n be arbitrary natural number. $(f_I v)^* = (f_{i_1^1} \cdots f_{i_1^1} v_1)^* \otimes \cdots \otimes (f_{i_n^n} \cdots f_{i_n^n} v_n)^*$. Formula (21) implies $\langle -x \cdot (f_I v)^*, \cdot \rangle = \langle (f_I v)^*, x \cdot \rangle = \langle (f_I v)^*, \sum_{j=1}^n x^{(j)} \cdot \rangle$. This and the computation for $n = 1$ give

$$-\Delta_{-, \alpha}(f_I v)^* = \sum_{(i_1, \dots, i_{m'}) \in P(\lambda', 1)} \sum_{j=1}^n \Delta_{i_1, \dots, i_{m'}} e_{i_{m'}}^{(j)} \cdots e_{i_1}^{(j)} (f_I v)^*. \quad \square$$

For every positive root $\alpha = \sum m_i^\alpha \alpha_i$ of \mathfrak{g} , set $\lambda_\alpha = (m_1^\alpha, \dots, m_r^\alpha) \in \mathbb{N}^r$. Now we combine Corollary 4.5 and Lemma 4.6 to obtain the following form of the Dynamical KZ equation.

Lemma 4.7 *Let $u(\mu, z) = \sum_{I \in P(\lambda, n)} u_I f_I v$, and let $\mu' \in \mathfrak{h}$ be a direction of differentiation. The Dynamical differential equation (13) is equivalent to the equation*

$$\begin{aligned} \kappa \partial_{\mu'} u &= \sum_{i=1}^n z_i (\mu')^{(i)} u + \\ &+ \sum_{I \in P(\lambda, n)} \sum_{\alpha > 0} \frac{\langle \alpha, \mu' \rangle}{\langle \alpha, \mu \rangle} \sum_{J \in P(\lambda, n)} \left\langle \left(\sum_{(i_1, \dots, i_{m'}) \in P(\lambda_\alpha, 1)} \Delta_{i_1, \dots, i_{m'}}(e_{i_{m'}} \cdots e_{i_1}) \right) (f_J v)^*, f_I v \right\rangle u_I f_J v. \quad \square \end{aligned} \quad (25)$$

4.4 A symmetrization procedure

The definition of the hypergeometric differential form involves a symmetrization procedure, see Section 3. Now we will study the behavior of the operator $\Delta_{i_1, \dots, i_{m'}} e_{i_{m'}} \cdots e_{i_1}$ for $(i_1, \dots, i_{m'}) \in P(\lambda', 1)$, where $\alpha = \alpha(\lambda')$, under the same type of symmetrization procedure.

Complexes [SV] For a Lie algebra \mathfrak{g} and a \mathfrak{g} -module M , denote by $C_\bullet(\mathfrak{g}, M)$ the standard chain complex of \mathfrak{g} with coefficients in M . $C_p(\mathfrak{g}, M) = \Lambda^p \mathfrak{g} \otimes M$ and

$$\begin{aligned} d : g_p \wedge \cdots \wedge g_1 \otimes x &= \sum_{i=1}^p (-1)^{i-1} g_p \wedge \cdots \wedge \widehat{g}_i \wedge \cdots \wedge g_1 \otimes g_i x + \\ &+ \sum_{1 \leq i < j \leq p} (-1)^{i+j} g_p \wedge \cdots \wedge \widehat{g}_j \wedge \cdots \wedge \widehat{g}_i \wedge \cdots \wedge g_1 \wedge [g_j, g_i] \otimes x. \end{aligned}$$

Let $\Lambda_1, \dots, \Lambda_n \in \mathfrak{h}^*$. Set $M = M(\Lambda_1) \otimes \cdots \otimes M(\Lambda_n)$. Consider the complex $C_\bullet(\mathfrak{n}_-, M)$. We have the weight decomposition $C_\bullet(\mathfrak{n}_-, M) = \bigoplus_{\lambda \in \mathbb{N}^r} C_\bullet(\mathfrak{n}_-, M)_\lambda$.

In Section 4.1 we recalled a Lie algebra structure on \mathfrak{n}_-^* and a \mathfrak{n}_-^* -module structure on M^* . Let $C_\bullet(\mathfrak{n}_-^*, M^*)$ be the corresponding standard chain complex: $C_\bullet(\mathfrak{n}_-^*, M^*) = \bigoplus_{\lambda \in \mathbb{N}^r} C_\bullet(\mathfrak{n}_-^*, M^*)_\lambda$. The covariant form induces a graded homomorphism of complexes, $S : C_\bullet(\mathfrak{n}_-, M) \rightarrow C_\bullet(\mathfrak{n}_-^*, M^*)$, see [SV].

Let $\lambda = (m_1, \dots, m_r) \in \mathbb{N}^r$, and $m = \sum m_i$. Define a subgroup Σ_λ of the symmetric group Σ_m via the direct product $\Sigma_\lambda = \Sigma_{m_1} \times \cdots \times \Sigma_{m_r}$, where Σ_{m_j} permutes the set of indices $\{\sum_{p=1}^{j-1} m_p + 1, \dots, \sum_{p=1}^j m_p + m_j\}$. Introduce a free Lie algebra $\widetilde{\mathfrak{n}}_-$ on generators $\widetilde{f}_1, \dots, \widetilde{f}_m$. Define a map of Lie algebras $\mathfrak{n}_- \rightarrow \widetilde{\mathfrak{n}}_-$ by setting $f_i \mapsto \sum_{j=1}^{m_i} \widetilde{f}_{m(i)+j}$, where $m(i) = m_1 + \cdots + m_{i-1}$. It induces a map of complexes $C_\bullet(\mathfrak{n}_-, U(\mathfrak{n}_-)^{\otimes n}) \rightarrow C_\bullet(\widetilde{\mathfrak{n}}_-, U(\widetilde{\mathfrak{n}}_-)^{\otimes n})^{\Sigma_\lambda}$. Set $\widetilde{\lambda} = \underbrace{(1, 1, \dots, 1)}_m$. Let

$$s : C_\bullet(\mathfrak{n}_-, U(\mathfrak{n}_-)^{\otimes n})_\lambda \rightarrow C_\bullet(\widetilde{\mathfrak{n}}_-, U(\widetilde{\mathfrak{n}}_-)^{\otimes n})_{\widetilde{\lambda}}^{\Sigma_\lambda} \quad (26)$$

be the previous map composed with the projection on the $\widetilde{\lambda}$ -component.

On the other hand there is a map of Lie algebras $\widetilde{\mathfrak{n}}_- \rightarrow \mathfrak{n}_-$ defined by $\widetilde{f}_j \mapsto f_i$, for $m(i) < j \leq m(i+1)$. It induces the map $\pi_\lambda : C_\bullet(\widetilde{\mathfrak{n}}_-, U(\widetilde{\mathfrak{n}}_-)^{\otimes n})_{\widetilde{\lambda}} \rightarrow C_\bullet(\mathfrak{n}_-, U(\mathfrak{n}_-)^{\otimes n})_\lambda$. Note that $s(y)$ equals the sum over the preimages of y under π_λ , for any $y \in C_\bullet(\mathfrak{n}_-, U(\mathfrak{n}_-)^{\otimes n})_\lambda$. Each such preimage is uniquely described by an element $\sigma \in \Sigma_\lambda$.

Example. Let $n = 1$ and $I = (i_1, \dots, i_m) \in P(\lambda, 1)$. Consider f_I as an element of $C_0(\mathfrak{n}_-, U(\mathfrak{n}_-))_\lambda$. Then $s(f_I) = \sum_{\sigma} \widetilde{f}_{\sigma_1} \cdots \widetilde{f}_{\sigma_m}$, where the sum is over the set $\{\sigma \in \Sigma(I)\} \cong \Sigma_\lambda$.

Lemma 4.8 *Let $n = 1$, $I \in P(\lambda, 1)$, and $m' \leq m$. Then the map $s^* : (C_1(\widetilde{\mathfrak{n}}_-, U(\widetilde{\mathfrak{n}}_-))_{\widetilde{\lambda}}^{\Sigma_\lambda})^* \rightarrow (C_1(\mathfrak{n}_-, U(\mathfrak{n}_-))_\lambda)^*$ has the following property*

$$s^* \left(\frac{1}{|\Sigma_\lambda|} \sum_{\sigma \in \Sigma(I)} \widetilde{\Delta}_{\sigma_1, \dots, \sigma_{m'}} \otimes (\widetilde{f}_{\sigma_{m'+1}} \cdots \widetilde{f}_{\sigma_m})^* \right) = \Delta_{i_1, \dots, i_{m'}} \otimes (f_{i_{m'+1}} \cdots f_{i_m})^* \quad (27)$$

Proof. Choose a basis (x_j^*) of \mathfrak{n}_-^* such that $x_1^* = \Delta_{i_1, \dots, i_{m'}}$ and let (x_j) be the dual basis of \mathfrak{n}_- . Take a tensor product of the basis (x_j) with the monomial basis in $U(\mathfrak{n}_-)$ to get a basis in $\mathfrak{n}_- \otimes U(\mathfrak{n}_-)$. Let us compare the two sides of (27) on that basis. $\Delta_{i_1, \dots, i_{m'}} \otimes (f_{i_{m'+1}} \cdots f_{i_m})^*(x_1 \otimes f_{i_{m'+1}} \cdots f_{i_m}) = 1$. The right hand side is zero on any other element of the basis.

Let $i : \mathfrak{n}_- \otimes U(\mathfrak{n}_-) \rightarrow U(\mathfrak{n}_-) \otimes U(\mathfrak{n}_-)$, and $\tilde{i} : \widetilde{\mathfrak{n}}_- \otimes U(\widetilde{\mathfrak{n}}_-) \rightarrow U(\widetilde{\mathfrak{n}}_-) \otimes U(\widetilde{\mathfrak{n}}_-)$ be the natural inclusion maps. Clearly $\tilde{i} \circ s = s \circ i$. Moreover, $\langle \tilde{\Delta}_{\sigma_1, \dots, \sigma_{m'}}, s(x) \rangle = \langle (\tilde{f}_{\sigma_1} \cdots \tilde{f}_{\sigma_{m'}})^*, (\tilde{i} \circ s(x)) \rangle$ by definition for $x \in \mathfrak{n}_-$. For a fixed $\sigma \in \Sigma_\lambda$ we have

$$\begin{aligned} & \langle s^*(\tilde{\Delta}_{\sigma_1, \dots, \sigma_{m'}} \otimes (\tilde{f}_{\sigma_{m'+1}} \cdots \tilde{f}_{\sigma_m})^*), x_1 \otimes f_{i_{m'+1}} \cdots f_{i_m} \rangle = \\ & = \langle \tilde{\Delta}_{\sigma_1, \dots, \sigma_{m'}} \otimes (\tilde{f}_{\sigma_{m'+1}} \cdots \tilde{f}_{\sigma_m})^*, s(x_1 \otimes f_{i_{m'+1}} \cdots f_{i_m}) \rangle \\ & = \langle (\tilde{f}_{\sigma_1} \cdots \tilde{f}_{\sigma_{m'}})^* \circ \tilde{i} \otimes (\tilde{f}_{\sigma_{m'+1}} \cdots \tilde{f}_{\sigma_m})^*, s(x_1 \otimes \tilde{f}_{i_{m'+1}} \cdots \tilde{f}_{i_m}) \rangle \\ & = \langle (\tilde{f}_{\sigma_1} \cdots \tilde{f}_{\sigma_{m'}})^* \otimes (\tilde{f}_{\sigma_{m'+1}} \cdots \tilde{f}_{\sigma_m})^*, s(i(x_1) \otimes \tilde{f}_{i_{m'+1}} \cdots \tilde{f}_{i_m}) \rangle \end{aligned}$$

The duality of the bases implies $i(x_1 \otimes f_{i_{m'+1}} \cdots f_{i_m}) = (f_{i_1} \cdots f_{i_{m'}} + \text{other monomials}) \otimes f_{i_{m'+1}} \cdots f_{i_m}$, and $s \circ i(x_1 \otimes f_{i_{m'+1}} \cdots f_{i_m}) = (\tilde{f}_{\sigma_1} \cdots \tilde{f}_{\sigma_{m'}} + \text{other monomials}) \otimes \tilde{f}_{\sigma_{m'+1}} \cdots \tilde{f}_{\sigma_m}$. Therefore, $\langle s^*(\tilde{\Delta}_{\sigma_1, \dots, \sigma_{m'}} \otimes (\tilde{f}_{\sigma_{m'+1}} \cdots \tilde{f}_{\sigma_m})^*), x_1 \otimes f_{i_{m'+1}} \cdots f_{i_m} \rangle = 1$. The same way we check that the left hand side is zero on the other basis elements. \square

Corollary 4.9 *Let π be the restriction of the projection π_λ to the subspace $C(\widetilde{\mathfrak{n}}_-, U(\widetilde{\mathfrak{n}}_-)^{\otimes n})_{\tilde{\lambda}}^{\Sigma_\lambda}$ of $C(\widetilde{\mathfrak{n}}_-, U(\widetilde{\mathfrak{n}}_-)^{\otimes n})_{\tilde{\lambda}}$. Let $J = (j_1, \dots, j_m) \in P(\lambda, n)$ and $I = (i_1, \dots, i_{m'}) \in P(\lambda_\alpha, 1)$ for a positive root α . Then we have*

$$\left(\sum_{\tau \in \Sigma(I)} \tilde{\Delta}_{\tau_1, \dots, \tau_{m'}}(\tilde{e}_{\tau_{m'}} \cdots \tilde{e}_{\tau_1}) \right) \left(\sum_{\sigma \in \Sigma(J)} (\tilde{f}_{\sigma_1} \cdots \tilde{f}_{\sigma_m})^* \right) = \pi^* \left((\Delta_{i_1, \dots, i_{m'}}(e_{i_{m'}} \cdots e_{i_1})) (f_{j_1} \cdots f_{j_m})^* \right). \quad (28)$$

Proof. Assume $n = 1$. The general case follows from this one because the operators $(\tilde{e}_{\tau_{m'}} \cdots \tilde{e}_{\tau_1})$ and $(e_{i_{m'}} \cdots e_{i_1})$ act on one tensor factor at a time. Since $s \circ \pi = |\Sigma_\lambda| \mathbf{id}$ we have $\pi^* \circ s^* = |\Sigma_\lambda| \mathbf{id}$. s^* and π^* are maps of complexes, i.e. they commute with the corresponding differentials. Thus, Lemma 4.8 implies after applying differentials and taking π^* from both sides

$$\pi^*(\Delta_{j_1, \dots, j_{m'}}(f_{j_{m'+1}} \cdots f_{j_m})^*) = \sum_{\sigma \in \Sigma(J)} \tilde{\Delta}_{\sigma_1, \dots, \sigma_{m'}}(\tilde{f}_{\sigma_{m'+1}} \cdots \tilde{f}_{\sigma_m})^*.$$

The right hand side of our formula is non-zero if and only if $m' \leq m$ and $i_k = j_k$ for $1 \leq k \leq m'$. Thus we compute

$$\begin{aligned} & \pi^*(\Delta_{i_1, \dots, i_{m'}} e_{i_{m'}} \cdots e_{i_1} (f_{j_1} \cdots f_{j_m})^*) = \pi^*(\Delta_{j_1, \dots, j_{m'}}(f_{j_{m'+1}} \cdots f_{j_m})^*) \\ & = \sum_{\sigma \in \Sigma(J)} \tilde{\Delta}_{\sigma_1, \dots, \sigma_{m'}}(\tilde{f}_{\sigma_{m'+1}} \cdots \tilde{f}_{\sigma_m})^* = \sum_{\sigma \in \Sigma(J)} \tilde{\Delta}_{\sigma_1, \dots, \sigma_{m'}} \tilde{e}_{\sigma_{m'}} \cdots \tilde{e}_{\sigma_1} (\tilde{f}_{\sigma_1} \cdots \tilde{f}_{\sigma_m})^* \end{aligned}$$

Note that $(\sigma_1, \dots, \sigma_{m'}) \in \Sigma(I)$. For any other $\tau \in \Sigma(I)$, we have $\tilde{e}_{\tau_{m'}} \cdots \tilde{e}_{\tau_1} (\tilde{f}_{\sigma_1} \cdots \tilde{f}_{\sigma_m})^* = 0$. Therefore we rewrite the last equality in the form (28). \square

Our last step is to show that the Dynamical equations for any Lie algebra in any weight space follow from the Dynamical equations for any Lie algebra in a weight space with weight $\lambda = (1, 1, \dots, 1)$.

Fix a finite dimensional complex vector space \mathbf{h} , a non-degenerate symmetric bilinear form (\cdot, \cdot) on \mathbf{h} , and a set of linearly independent ‘‘simple roots’’ $\alpha_1, \dots, \alpha_r \in \mathbf{h}^*$. Consider the corresponding Kac-Moody Lie algebra without Serre’s relations, \mathbf{g} , defined at the beginning of Section 4.1. Recall that $b : \mathbf{h} \rightarrow \mathbf{h}^*$ denotes the isomorphism induced by the bilinear form, $h_i = b^{-1}(\alpha_i)$, and the form (\cdot, \cdot) is transferred to \mathbf{h}^* via the map b .

Fix $\lambda = (m_1, \dots, m_r) \in \mathbb{N}^r$. Consider the corresponding positive root $\alpha(\lambda) = \sum_{i=1}^r m_i \alpha_i$ of \mathbf{g} . Up to reordering of the α ’s we can assume that $\alpha(\lambda) = \sum_{i=1}^p m_i \alpha_i$ where $m_i > 0$ for $1 \leq i \leq p$ and $p \leq r$ is fixed. The corresponding coloring function is $c_\lambda : \{1, \dots, m = \sum m_i\} \rightarrow \{1, \dots, p\}$. We use the following linear algebraic fact when symmetrizing.

Proposition 4.10 *Let \mathbf{h} be a finite dimensional vector space with a non-degenerate symmetric bilinear form (\cdot, \cdot) , and a set of linearly independent vectors $(h_i)_{i=1}^r \subset \mathbf{h}$. Then there exists a finite dimensional vector space $\tilde{\mathbf{h}}$ with a non-degenerate symmetric bilinear form $(\cdot, \cdot)_1$, a set of linearly independent vectors $(\tilde{h}_j)_{j=1}^m \subset \tilde{\mathbf{h}}$, and a monomorphism $s_h : \mathbf{h} \rightarrow \tilde{\mathbf{h}}$ such that*

- (a) $s_h(h_i) = \frac{1}{m_i} \sum_{j=1}^{m_i} \tilde{h}_{m(i)+j}$, where $m(i) = m_1 + \dots + m_{i-1}$, $i = 1, \dots, p$;
- (b) $(\tilde{h}_j, s_h(h'))_1 = (h_{c(j)}, h')$ and $(h', h'') = (s_h(h'), s_h(h''))_1$ for any $h', h'' \in \mathbf{h}$, $j = 1 \dots, m$.

Proof. Let $q = \dim \mathbf{h}$. Complete the set h_1, \dots, h_r to a basis $h_1, \dots, h_r, h_{r+1}, \dots, h_q$ of \mathbf{h} . U Consider a complex linear space $\tilde{\mathbf{h}}' = \mathbb{C}\{\tilde{h}_1, \dots, \tilde{h}_m, \tilde{h}_{m+1}, \dots, \tilde{h}_{m+q-p}\}$. Extend the coloring function $c : \{1, \dots, m+q-p\} \rightarrow \{1, \dots, q\}$ setting $c(m+j) = r+j$ for $j = 1, \dots, q-p$. Define a symmetric degenerate bilinear form on $\tilde{\mathbf{h}}'$ by the rules $(\tilde{h}_j, \tilde{h}_k)_1 = (h_{c(j)}, h_{c(k)})$ for $1 \leq j, k \leq m+q-p$. The rank of the form is q and the dimension of its kernel is $m-p$. There exists an extension $\tilde{\mathbf{h}}$ of the vector space $\tilde{\mathbf{h}}'$ and an extension of $(\cdot, \cdot)_1$ to a non-degenerate symmetric bilinear form on $\tilde{\mathbf{h}}$.

Define a monomorphism s_h by $s_h(h_i) = \frac{1}{m_i} \sum_{j=1}^{m_i} \tilde{h}_{m(i)+j}$, where $m(i) = m_1 + \dots + m_{i-1}$, $i = 1, \dots, q$, and note that $s_h(h_{p+j}) = \tilde{h}_{m+j}$ for $j = 1, \dots, q-p$. Now checking (b) on a basis is straightforward. \square

Set $\tilde{\alpha}_j = (\tilde{h}_j, \cdot)_1 \in \tilde{\mathbf{h}}^*$ for $j = 1 \dots, m$. Consider $\tilde{\mathbf{g}}$, a Kac-Moody Lie algebra without Serre’s relations corresponding to the data $\tilde{\mathbf{h}}$, $(\cdot, \cdot)_1$, and $(\tilde{\alpha}_j)_{j=1}^m$. Note that $1 \leq j \leq m$ implies $\langle \tilde{\alpha}_j, s_h(h') \rangle = (\tilde{h}_j, s_h(h'))_1 = (h_{c(j)}, h') = \langle \alpha_{c(j)}, h' \rangle$ for any $h' \in \mathbf{h}$.

Let $M = M(\Lambda_1) \otimes \dots \otimes M(\Lambda_n)$ be a tensor product of Verma modules for \mathbf{g} with corresponding highest weights $\Lambda_1, \dots, \Lambda_n \in \mathbf{h}^*$. Since s_h is a monomorphism, $s_h^* : \tilde{\mathbf{h}}^* \rightarrow \mathbf{h}^*$ is a linear epimorphism. Choose highest weights $\tilde{\Lambda}_1, \dots, \tilde{\Lambda}_n \in \tilde{\mathbf{h}}^*$ such that $s_h^*(\tilde{\Lambda}_j) = \Lambda_j$ for $1 \leq j \leq n$, and consider the corresponding tensor product of Verma modules for $\tilde{\mathbf{g}}$, $\tilde{M} = \tilde{M}(\tilde{\Lambda}_1) \otimes \dots \otimes \tilde{M}(\tilde{\Lambda}_n)$.

Lemma 4.11 Let $\tilde{\lambda} = \underbrace{(1, 1, \dots, 1)}_m$. Let $\tilde{u}(\tilde{\mu}, z) = \sum_{K \in P(\tilde{\lambda}, n)} \tilde{u}_K \tilde{f}_K$ be a hypergeometric solution of the Dynamical equations with values in the $\tilde{\lambda}$ weight space of a $\tilde{\mathfrak{g}}$ -module $\tilde{M} \cong U(\tilde{\mathfrak{n}}_-)^{\otimes n}$. Then $u(\mu, z) = \pi(\tilde{u})(s_h(\mu), z)$ is a hypergeometric solution of the Dynamical equations with values in the λ weight space of a \mathfrak{g} -module $M \cong U(\mathfrak{n}_-)^{\otimes n}$, i.e. $u = \sum_{I \in P(\lambda, n)} u_I f_I$.

Proof. Note that by definition $P(\tilde{\lambda}, n) = \cup_{I \in P(\lambda, n)} \{K \in \Sigma(I)\}$. From the definition of the hypergeometric differential form, see Section 3, it follows that $u_I = \sum_{k \in \Sigma(I)} \tilde{u}_k$. Therefore

$$\pi(\tilde{u}) = \pi\left(\sum_{I \in P(\lambda, n)} \sum_{K \in \Sigma(I)} \tilde{u}_K \tilde{f}_K\right) = \sum_{I \in P(\lambda, n)} \left(\sum_{K \in \Sigma(I)} \tilde{u}_K\right) f_I = \sum_{I \in P(\lambda, n)} u_I f_I. \quad (29)$$

Fix a point $\mu \in \mathfrak{h}$ and a direction of differentiation $\mu' \in \mathfrak{h}$. Denote $\tilde{\mu} = s_h(\mu)$ and $\tilde{\mu}' = s_h(\mu')$. Since

$$\begin{aligned} \partial_{\mu'} \exp\left(-\sum_{j=1}^m \langle \alpha_{c(j)}, \mu \rangle t_j + \sum_{l=1}^n \langle \Lambda_l, \mu \rangle\right) &= \\ &= \left(-\sum_{j=1}^m \langle \alpha_{c(j)}, \mu' \rangle t_j + \sum_{l=1}^n \langle \Lambda_l, \mu' \rangle\right) \exp\left(-\sum_{j=1}^m \langle \alpha_{c(j)}, \mu \rangle t_j + \sum_{l=1}^n \langle \Lambda_l, \mu \rangle\right) \\ &= \left(-\sum_{j=1}^m \langle \tilde{\alpha}_j, \tilde{\mu}' \rangle t_j + \sum_{l=1}^n \langle \tilde{\Lambda}_l, \tilde{\mu}' \rangle\right) \exp\left(-\sum_{j=1}^m \langle \tilde{\alpha}_j, \tilde{\mu} \rangle t_j + \sum_{l=1}^n \langle \tilde{\Lambda}_l, \tilde{\mu} \rangle\right) \\ &= \partial_{\tilde{\mu}'} \exp\left(-\sum_{j=1}^m \langle \tilde{\alpha}_j, \tilde{\mu} \rangle t_j + \sum_{l=1}^n \langle \tilde{\Lambda}_l, \tilde{\mu} \rangle\right), \end{aligned} \quad (30)$$

we have $\pi(\partial_{\tilde{\mu}'} \tilde{u}(\tilde{\mu}, z)) = \partial_{\mu'} \pi(\tilde{u})(\mu, z)$. If $I = (I_1, \dots, I_n) \in P(\lambda, n)$ and $K = (K_1, \dots, K_n) \in \Sigma(I)$, where $K_j = (k_1^j, \dots, k_{s_j}^j)$ and $I_j = (i_1^j, \dots, i_{s_j}^j)$, then

$$\begin{aligned} \tilde{\mu}'^{(j)} \tilde{f}_K &= \langle \tilde{\Lambda}_j - \sum_{l=1}^{s_j} \tilde{\alpha}_{k_l^j}, \tilde{\mu}' \rangle \tilde{f}_K = \langle \Lambda_j - \sum_{l=1}^{s_j} \alpha_{i_l^j}, \mu' \rangle \tilde{f}_K, \\ \pi(\tilde{\mu}'^{(j)} \sum_{K \in \Sigma(I)} \tilde{u}_K \tilde{f}_K) &= \langle \Lambda_j - \sum_{l=1}^{s_j} \alpha_{i_l^j}, \mu' \rangle \left(\sum_{K \in \Sigma(I)} \tilde{u}_K \pi(\tilde{f}_K)\right) \\ &= \langle \Lambda_j - \sum_{l=1}^{s_j} \alpha_{i_l^j}, \mu' \rangle \left(\sum_{K \in \Sigma(I)} \tilde{u}_K\right) f_I = \langle \Lambda_j - \sum_{l=1}^{s_j} \alpha_{i_l^j}, \mu' \rangle u_I f_I = \mu'^{(j)} f_I \end{aligned} \quad (31)$$

Combine formulae (29) and (31) to obtain

$$\pi\left(\sum_{j=1}^n z_j \tilde{\mu}'^{(j)} \tilde{u}\right) = \sum_{j=1}^n z_j \mu'^{(j)} u. \quad (32)$$

Let $\alpha = \sum_{i=1}^r m'_i \alpha_j$ be a positive root for \mathfrak{g} . Lemma 4.6 gives a necessary condition for a non-zero action of $\Delta_{-, \alpha}$ on M_λ^* . Namely $m'_i \leq m_i$ for all $i = 1, \dots, r$. Analogously, for a positive root $\tilde{\alpha} = \sum_{j=1}^m p_j \tilde{\alpha}_j$ of $\tilde{\mathfrak{g}}$ a necessary condition for a non-zero action of $\Delta_{-, \tilde{\alpha}}$ on \tilde{M}_λ^* is $p_j = 0, 1$ for $j = 1, \dots, m$. Call all such α 's ($\tilde{\alpha}$'s) λ -admissible ($\tilde{\lambda}$ -admissible). Since $s_h^*(\tilde{\alpha}_j) = \alpha_{c(j)}$, s_h^* maps the set of $\tilde{\lambda}$ -admissible roots of $\tilde{\mathfrak{g}}$ onto the set of λ -admissible roots of \mathfrak{g} .

Let $\alpha = \sum m'_i \alpha_i$ be a λ -admissible root for \mathfrak{g} and $m' = \sum m'_i$. For any $\tilde{\alpha}$, such that $s_h^*(\tilde{\alpha}) = \alpha$, we have $\langle \tilde{\alpha}, \tilde{\mu}' \rangle / \langle \tilde{\alpha}, \tilde{\mu} \rangle = \langle \alpha, \mu' \rangle / \langle \alpha, \mu \rangle$. Consider $\pi(\sum_{\tilde{\alpha}, s_h^*(\tilde{\alpha})=\alpha} \frac{\langle \tilde{\alpha}, \tilde{\mu}' \rangle}{\langle \tilde{\alpha}, \tilde{\mu} \rangle} \Delta_{+, \tilde{\alpha}} \tilde{u})$, where the sum is over $\tilde{\lambda}$ -admissible roots. Corollary 4.5 applied to the basis $(\tilde{f}_K)_{K \in P(\tilde{\lambda}, n)}$ of M_λ gives

$$\begin{aligned} & \frac{\langle \alpha, \mu \rangle}{\langle \alpha, \mu' \rangle} \pi \left(\sum_{\tilde{\alpha}, s_h^*(\tilde{\alpha})=\alpha} \frac{\langle \tilde{\alpha}, \tilde{\mu}' \rangle}{\langle \tilde{\alpha}, \tilde{\mu} \rangle} \Delta_{+, \tilde{\alpha}} \tilde{u} \right) = \pi \left(\sum_{K \in P(\tilde{\lambda}, n)} \left\langle - \sum_{\tilde{\alpha}, s_h^*(\tilde{\alpha})=\alpha} \Delta_{-, \tilde{\alpha}} (\tilde{f}_K)^*, \tilde{u} \right\rangle \tilde{f}_K \right) = \\ & = \sum_{I \in P(\lambda, n)} \left\langle \left(- \sum_{\tilde{\alpha}, s_h^*(\tilde{\alpha})=\alpha} \Delta_{-, \tilde{\alpha}} \right) \left(\sum_{K \in \Sigma(I)} (\tilde{f}_K)^* \right), \tilde{u} \right\rangle f_I \end{aligned} \quad (33)$$

Lemma 4.6 asserts that

$$- \sum_{\tilde{\alpha}, s_h^*(\tilde{\alpha})=\alpha} \Delta_{-, \tilde{\alpha}} = \sum_{\tilde{\alpha}, s_h^*(\tilde{\alpha})=\alpha} \left(\sum_{(l_1, \dots, l_{m'}) \in P(\tilde{\lambda}_{\tilde{\alpha}}, 1)} \tilde{\Delta}_{l_1, \dots, l_{m'}} (\tilde{e}_{l_{m'}} \dots \tilde{e}_{l_1}) \right).$$

Rearrange the summation using that sum over $(l_1, \dots, l_{m'}) \in P(\tilde{\lambda}_{\tilde{\alpha}}, 1)$ such that $s_h^*(\tilde{\alpha}) = \alpha$ equals the sum over $(p_1, \dots, p_{m'}) \in \Sigma(J)$ such that $J = (j_1, \dots, j_{m'}) \in P(\lambda_\alpha, 1)$. Combine such rearrangement with Lemma 4.6 and Corollary 4.9 to simplify formula (33).

$$\begin{aligned} & \frac{\langle \alpha, \mu \rangle}{\langle \alpha, \mu' \rangle} \pi \left(\sum_{\tilde{\alpha}, s_h^*(\tilde{\alpha})=\alpha} \frac{\langle \tilde{\alpha}, \tilde{\mu}' \rangle}{\langle \tilde{\alpha}, \tilde{\mu} \rangle} \Delta_{+, \tilde{\alpha}} \tilde{u} \right) = \\ & = \sum_{I \in P(\lambda, n)} \left\langle \sum_{J \in P(\lambda_\alpha, 1)} \left(\sum_{(p_1, \dots, p_{m'}) \in \Sigma(J)} \tilde{\Delta}_{p_1, \dots, p_{m'}} (\tilde{e}_{p_{m'}} \dots \tilde{e}_{p_1}) \right) \left(\sum_{K \in \Sigma(I)} (\tilde{f}_K)^* \right), \tilde{u} \right\rangle f_I \\ & = \sum_{I \in P(\lambda, n)} \left\langle \sum_{J \in P(\lambda_\alpha, 1)} \pi^* (\Delta_{j_1, \dots, j_{m'}} (e_{j_{m'}} \dots e_{j_1}) (f_I)^*), \tilde{u} \right\rangle f_I \\ & = \sum_{I \in P(\lambda, n)} \left\langle \sum_{J \in P(\lambda_\alpha, 1)} \Delta_{j_1, \dots, j_{m'}} (e_{j_{m'}} \dots e_{j_1}) (f_I)^*, \pi(\tilde{u}) \right\rangle f_I \\ & = \sum_{I \in P(\lambda, n)} \langle -\Delta_{-, \alpha} (f_I)^*, u \rangle f_I = \Delta_{+, \alpha} u. \end{aligned} \quad (34)$$

Finally (32) and (34) imply

$$\partial_{\mu'} u = \pi(\partial_{\tilde{\mu}'} \tilde{u}) = \pi \left(\left(\sum_{j=1}^n z_j \tilde{\mu}'^{(j)} + \sum_{\tilde{\alpha} > 0} \frac{\langle \tilde{\alpha}, \tilde{\mu}' \rangle}{\langle \tilde{\alpha}, \tilde{\mu} \rangle} \Delta_{+, \tilde{\alpha}} \tilde{u} \right) \right) = \left(\sum_{j=1}^n z_j \mu'^{(j)} + \sum_{\alpha > 0} \frac{\langle \alpha, \mu' \rangle}{\langle \alpha, \mu \rangle} \Delta_{+, \alpha} \right) u. \quad \square \quad (35)$$

4.5 The proof of Lemma 4.1

Recall that the linear map $\nu_M : M \rightarrow \mathfrak{b}_- \otimes M$ has the following property $\nu_M(x) = \frac{1}{2}(b^{-1}(\Lambda - \alpha(\lambda))) \otimes x + \nu_{M^-}(x)$, where $x \in M_\lambda$, $\nu_{M^-}(x) \in \mathfrak{n}_- \otimes M$, and $b^{-1} : \mathfrak{h}^* \rightarrow \mathfrak{h}$ is defined at the beginning of Section 4.1. Let $a \in \mathfrak{h}$, $x \in M_\lambda$. Since $S(b \otimes x, a \otimes y) = S(a, b)S(x, y)$ for any $b \in \mathfrak{g}$, $y \in M$, and \mathfrak{h} is orthogonal to \mathfrak{n}_- with respect to S , and (\cdot, \cdot) coincides with S on \mathfrak{h} we have

$$S(\nu_{M^-}(x), a \otimes y) = S\left(\frac{1}{2}(b^{-1}(\Lambda - \alpha(\lambda))), a\right)S(x, y) = \frac{1}{2}S(x, ay). \quad (36)$$

This proves the first equality in the Lemma. To prove the second part for a monomial x , we use double induction by the number of tensor factors and the number of f 's in x . We use $\nu_{M^-}(x)$ instead of $\nu_M(x)$ because of the orthogonality mentioned above.

Let $M = M(\Lambda_1)$ be a highest weight module of \mathfrak{g} with a highest vector v , and $a \in \mathfrak{n}_-$, $y \in M$. Since $S(\nu_{M^-}(v)) = 0$, we have $S(\nu_{M^-}(v), a \otimes y) = 0 = S(v, ay)$. The inductive step is as follows. Assume $S(\nu_{M^-}(x), a \otimes y) = S(x, ay)$. Then

$$\begin{aligned} S(\nu_{M^-}(f_i x), a \otimes y) &= \\ &= S(f_i \otimes h_i x, a \otimes y) + S(f_i \nu_{M^-}(x), a \otimes y) = S(f_i, a)S(h_i x, y) + S(\nu_{M^-}(x), e_i(a \otimes y)) \\ &= S(f_i, a)S(h_i x, y) + S(\nu_{M^-}(x), [e_i, a] \otimes y) + S(\nu_{M^-}(x), a \otimes e_i y) \\ &= S(f_i, a)S(h_i x, y) + S(\nu_{M^-}(x), [e_i, a] \otimes y) + S(x, a e_i y) \\ &= S(f_i, a)S(h_i x, y) + S(\nu_{M^-}(x), [e_i, a] \otimes y) + S(x, e_i a y) - S(x, [e_i, a] y) \\ &= S(f_i, a)S(h_i x, y) + S(\nu_{M^-}(x), [e_i, a] \otimes y) + S(f_i x, a y) - S(x, [e_i, a] y) \end{aligned} \quad (37)$$

If $a = f_i$, then (37) and the properties $S(x, h y) = S(h x, y)$, $S(\nu_{M^-}(x), h \otimes y) = 0$ for $h \in \mathfrak{h}$ imply

$$S(\nu_{M^-}(f_i x), a \otimes y) = S(h_i x, y) + S(\nu_{M^-}(x), h_i \otimes y) + S(f_i x, a y) - S(x, h_i y) = S(f_i x, y).$$

If a is orthogonal to f_i with respect to S , then (37) and the inductive hypothesis give

$$S(\nu_{M^-}(f_i x), a \otimes y) = 0 + S(x, [e_i, a] y) + S(f_i x, a y) - S(x, [e_i, a] y) = S(f_i x, a).$$

Thus the statement is proved for one tensor factor.

Assume that $S(\nu_{M^-}(f_i x), a \otimes y) = S(x, a y)$ for a module M , which is a tensor product of up to $n - 1$ tensor factors ($n \geq 2$).

Let $M = M(\Lambda_1) \otimes \cdots \otimes M(\Lambda_n)$. Recall that $\nu_{M^-}(x) = \sum_{k=1}^n \nu_M^{(k)}(x)_-$, where $\nu_M^{(k)}(f_i^{(j)} x)_- = f_i^{(j)} \nu_M^{(k)}(x)_-$ for $k \neq j$, and $\nu_M^{(k)}(f_i^{(k)} x)_- = f_i \otimes h_i^{(k)} x + f_i^{(k)} \nu_M^{(k)}(x)_-$. The following commutation relations will be useful. $S(f_i^{(j)} \nu_{M^-}^{(k)}(x), a \otimes y) = S(\nu_{M^-}^{(k)}(x), a \otimes e_i^{(j)} y)$, for $j \neq k$, and $S(f_i^{(j)} \nu_{M^-}^{(j)}(x), a \otimes y) = S(\nu_{M^-}^{(j)}(x), [e_i, a] \otimes y) + S(\nu_{M^-}^{(j)}(x), a \otimes e_i^{(j)} y)$. Both equalities are corollaries of the Lemma 4.1 for one tensor factor, and the definition of S , e.g. $S(x_1 \otimes \cdots \otimes x_{j-1} \otimes \nu_{M^-}(x_j) \otimes x_{j+1} \otimes \cdots \otimes x_n, a \otimes y_1 \otimes \cdots \otimes y_j \otimes \cdots \otimes y_n) = S(\nu_{M^-}(x_j), a \otimes y_j) \prod_{k \neq j} S(x_k, y_k)$.

In all formulae $1 \leq i \leq r$, and $1 \leq j, k \leq n$, and the upper script indicates the tensor factor where the action is applied. Let $a \in \mathfrak{n}_-$. The base for the induction is exactly as for $n = 1$. The inductive step is as follows.

$$\begin{aligned}
S(\nu_{M^-}(f_i^{(j)}x), a \otimes y) &= \sum_{k \neq j} S(\nu_{M^-}^{(k)}(f_i^{(j)}x), a \otimes y) + S(\nu_{M^-}^{(j)}(f_i^{(j)}x), a \otimes y) \\
&= \sum_{k \neq j} S(f_i^{(j)}\nu_{M^-}^{(k)}(x), a \otimes y) + S(f_i \otimes h_i^{(j)}x, a \otimes y) + S(f_i^{(j)}\nu_M^{(j)}(x)_-, a \otimes y) \\
&= S(f_i, a)S(h_i^{(j)}x, y) + S(\nu_{M^-}^{(j)}(x), [e_i, a] \otimes y) + \sum_{k=1}^n S(\nu_{M^-}^{(k)}(x), a \otimes e_i^{(j)}y). \tag{38}
\end{aligned}$$

The result for one tensor factor gives

$$\begin{aligned}
\sum_k S(\nu_{M^-}^{(k)}(x), a \otimes e_i^{(j)}y) &= \sum_k S(x, a^{(k)}e_i^{(j)}y) = \sum_k S(x, e_i^{(j)}a^{(k)}y) - S(x, [e_i, a]^{(j)}y) \\
&= S(f_i^{(j)}x, ay) - S(x, [e_i, a]^{(j)}y). \tag{39}
\end{aligned}$$

If $a = f_i$, then (38), (39) and the properties $S(x, h^{(k)}y) = S(h^{(k)}x, y)$, $S(\nu_{M^-}^{(k)}(x), h \otimes y) = 0$ for $h \in \mathfrak{h}$, $k = 1, \dots, n$ imply

$$S(\nu_{M^-}(f_i^{(j)}x), a \otimes y) = S(h_i^{(j)}x, y) + 0 + S(f_i^{(j)}x, ay) - S(x, h_i^{(j)}y) = S(f_i^{(j)}x, ay).$$

If a is orthogonal to f_i with respect to S , then (38), (39) and the result for one tensor factor give

$$S(\nu_{M^-}(f_i^{(j)}x), a \otimes y) = 0 + S(x, [e_i, a]^{(j)}y) + S(f_i^{(j)}x, ay) - S(x, [e_i, a]^{(j)}y) = S(f_i^{(j)}x, ay).$$

This finishes the inductive argument. The Lemma is proved. \square

5 Flags, Orlik-Solomon algebra, hypergeometric differential forms

In this section we will formulate results from [SV] which define a map between the complex of hypergeometric differential forms and the complex $C_\bullet(\mathfrak{n}_-^*, M^*)$ for a suitable Lie algebra \mathfrak{n}_- and a \mathfrak{n}_- -module M .

5.1 Complexes

Let W be an affine complex m -dimensional space and let \mathcal{C} be a configuration of hyperplanes in W . Define Abelian groups $\mathcal{A}^k(\mathcal{C}, \mathbb{Z})$, $0 \leq k \leq m$, as follows. $\mathcal{A}^0(\mathcal{C}, \mathbb{Z}) = \mathbb{Z}$. For $k \geq 1$, $\mathcal{A}^k(\mathcal{C})$ is generated by k -tuples (H_1, \dots, H_k) , $H_i \in \mathcal{C}$, subject to the relations:

$$(H_1, \dots, H_k) = 0 \text{ if } H_1, \dots, H_k \text{ are not in general position (codim } H_1 \cap \dots \cap H_k \neq k);$$

$(H_{\sigma(1)}, \dots, H_{\sigma(k)}) = (-1)^{|\sigma|} (H_1, \dots, H_k)$ for any permutation $\sigma \in \Sigma_k$;

$\sum_{i=1}^{k+1} (-1)^i (H_1, \dots, \hat{H}_i, \dots, H_{k+1}) = 0$ for any $(k+1)$ -tuple H_1, \dots, H_{k+1} which is not in general position and such that $H_1 \cap \dots \cap H_k \neq 0$.

The direct sum $\mathcal{A}^\bullet(\mathcal{C}, \mathbb{Z}) = \bigoplus_{k=0}^m \mathcal{A}^k(\mathcal{C}, \mathbb{Z})$ is a graded skew commutative algebra with respect to the multiplication $(H_1, \dots, H_k) \cdot (H'_1, \dots, H'_l) = (H_1, \dots, H_k, H'_1, \dots, H'_l)$. $\mathcal{A}^\bullet(\mathcal{C}, \mathbb{Z})$ is called the *Orlik-Solomon algebra* of the configuration \mathcal{C} .

Flags. For $0 \leq k \leq m$, denote by $\text{Flag}^k(\mathcal{C})$ the set of all flags $L^0 \supset L^1 \supset \dots \supset L^k$, where L^i is an edge of \mathcal{C} of codimension i . Denote by $\overline{\text{Flag}}^k(\mathcal{C})$ the free Abelian group on $\text{Flag}^k(\mathcal{C})$ and by $Fl^k(\mathcal{C}, \mathbb{Z})$ the quotient of $\overline{\text{Flag}}^k(\mathcal{C})$ by the following relations.

For every i , $0 < i < k$, and a flag with a gap, $\hat{F} = (L^0 \supset \dots \supset L^{i-1} \supset L^{i+1} \supset L^k)$, where L^j is an edge of codimension j , we set $\sum_{F \supset \hat{F}} F = 0$ in $Fl^k(\mathcal{C}, \mathbb{Z})$, where the summation is over all flags $F = (\tilde{L}^0 \supset \tilde{L}^k) \in \text{Flag}^k(\mathcal{C})$ such that $\tilde{L}^j = L^j$ for all $j \neq i$.

To define the relation between $\mathcal{A}^k(\mathcal{C}, \mathbb{Z})$ and $Fl^k(\mathcal{C}, \mathbb{Z})$ we define the following map. For (H_1, \dots, H_k) in the general position, $H_i \in \mathcal{C}$, define $F(H_1, \dots, H_k) = (H_1 \supset H_{12} \supset \dots \supset H_{12\dots k}) \in \text{Flag}^k(\mathcal{C})$, where $H_{12\dots i} = H_1 \cap H_2 \cap \dots \cap H_i$. For a flag $F \in \text{Flag}^k(\mathcal{C})$, define a functional $\delta_F \in Fl^k(\mathcal{C}, \mathbb{Z})^*$ as $\delta_F(F') = 1$ if $F' = F$ and $\delta_F(F') = 0$ otherwise. For (H_1, \dots, H_k) in general position, define a map

$$\varphi^k(H_1, \dots, H_k) = \sum_{\sigma \in \Sigma_k} (-1)^{|\sigma|} \delta_{F(H_{\sigma_1}, \dots, H_{\sigma_k})}. \quad (40)$$

Thus we have a homomorphism $\varphi^k : \mathcal{A}^k(\mathcal{C}, \mathbb{Z}) \rightarrow Fl^k(\mathcal{C}, \mathbb{Z})^*$. The following statements are from [SV]. All groups $Fl^k(\mathcal{C}, \mathbb{Z})$ are free over \mathbb{Z} . $\mathcal{A}^k(\mathcal{C}, \mathbb{Z})$ and $Fl^k(\mathcal{C}, \mathbb{Z})$ are dual and the map φ^k is an isomorphism.

Set $\mathcal{A}^k(\mathcal{C}) = \mathcal{A}^k(\mathcal{C}, \mathbb{Z}) \otimes_{\mathbb{Z}} \mathbb{C}$ and $Fl^k(\mathcal{C}) = Fl^k(\mathcal{C}, \mathbb{Z}) \otimes_{\mathbb{Z}} \mathbb{C}$ for all k .

From now on we assume that the configuration \mathcal{C} is weighted, that is, to any hyperplane $H \in \mathcal{C}$ its weight, a number $a(H) \in \mathbb{C}$, is assigned. Define the quasiclassical weight of any edge L of \mathcal{C} as the sum of the weights of all hyperplanes that contain the edge.

Say that a k -tuple $\bar{H} = (H_1, \dots, H_k)$, $H_i \in \mathcal{C}$, is adjacent to a flag F if there exists $\sigma \in \Sigma_k$ such that $F = F(H_{\sigma_1}, \dots, H_{\sigma_k})$. This permutation σ is unique. Denote it by $\sigma(\bar{H}, F)$.

Define a symmetric bilinear form S^k on $Fl^k(\mathcal{C})$. For $F, F' \in \text{Flag}^k(\mathcal{C})$, set

$$S^k(F, F') = \frac{1}{k} \sum (-1)^{\sigma(\bar{H}, F)\sigma(\bar{H}, F')} a(H_1) \dots a(H_k), \quad (41)$$

where the summation is over all $\bar{H} = (H_1, \dots, H_k)$ adjacent to both F and F' .

The form S^k is called the *quasiclassical contravariant form* of the configuration \mathcal{C} . It defines a bilinear symmetric form on $Fl^k(\mathcal{C})$. See [SV].

Flag complex. Define a differential $d : Fl^k \rightarrow Fl^{k+1}$ by $d(L^0 \supset \dots \supset L^k) = \sum_{L^{k+1}} (L^0 \supset \dots \supset L^k \supset L^{k+1})$, where the sum is taken over all edges L^{k+1} of codimension $k+1$ such that $L^k \supset L^{k+1}$. From the definition of the groups Fl^k it follows that $d^2 = 0$.

A complex $(\mathcal{A}^\bullet, d(a))$. Set $\omega = \omega(a) = \sum_{H \in \mathcal{C}} a(H)H$, $\omega(a) \in \mathcal{A}^1$. Define a differential $d = d(a) : \mathcal{A}^k \rightarrow \mathcal{A}^{k+1}$ by the rule $dx = \omega(a) \cdot x$. It is clear that $d^2 = 0$.

For any k , the quasiclassical bilinear form on \mathcal{C} defines a homomorphism

$$S^k : Fl^k \rightarrow (Fl^k)^* \simeq \mathcal{A}^k, \quad (42)$$

where $S^k(F) = (-1)^{k(k-1)/2} S(F, \cdot)$.

Lemma 5.1 S^\bullet defines a map of complexes $S^\bullet = S^\bullet(a) : (Fl^\bullet(\mathcal{C}), d) \rightarrow (\mathcal{A}^\bullet(\mathcal{C}), d(a))$.

Note. There is a misprint in [SV] in the definition of S^k where the factor $(-1)^{k(k-1)/2}$ is missing.

Proof. For any edge L , set $S(L) = \sum_{H \in \mathcal{C}, L \subset H} a(H)H$, $S(L) \in \mathcal{A}^1$. It is easy to see that the homomorphism S^k is defined by $S^k(L^0 \supset \dots \supset L^k) = (-1)^{k(k-1)/2} S(L^1) \cdot S(L^2) \dots S(L^k)$. In other words $S^k(L^0 \supset \dots \supset L^k) = (-1)^{k(k-1)/2} \sum a(H_1) \dots a(H_k)(H_1, \dots, H_k)$, where the sum is over all k -tuples (H_1, \dots, H_k) such that $H_i \supset L^i$ for all i . Therefore, we have

$$\begin{aligned} S^{k+1}d(L^0 \supset \dots \supset L^k) &= S^{k+1} \left(\sum_{L^{k+1}, L^{k+1} \subset L^k} (L^0 \supset \dots \supset L^k \supset L^{k+1}) \right) \\ &= (-1)^{\frac{(k+1)k}{2}} \sum a(H_1) \dots a(H_k) a(H_{k+1})(H_1, \dots, H_k, H_{k+1}) \\ &= \left((-1)^{\frac{k(k-1)}{2}} \sum a(H_1) \dots a(H_k)(H_1, \dots, H_k) \right) \cdot (-1)^k \sum a(H_{k+1})H_{k+1} \\ &= \left((-1)^{\frac{k(k-1)}{2}} \sum a(H_1) \dots a(H_k)(H_1, \dots, H_k) \right) \cdot (-1)^k \omega(a) \\ &= S^k(L^0 \supset \dots \supset L^k) \cdot (-1)^k \omega(a) = \omega(a) \cdot S^k(L^0 \supset \dots \supset L^k) = d(a)S^k(L^0 \supset \dots \supset L^k). \end{aligned}$$

The second, third and fourth sum are over all H_i , such that $H_i \supset L^i$, for $1 \leq i \leq k$ and $H_{k+1} \cap L^k \neq 0$. Note that $H_{k+1} \cap L^k = 0$ implies $(H_1, \dots, H_k, H_{k+1}) = 0$, and thus the fourth equality is justified. The sixth one comes from the skew symmetry in $\mathcal{A}^\bullet(\mathcal{C})$. \square

Recall that we have a weighted configuration of hyperplanes in a complex m - dimensional space W , and $a = \{a(H) | H \in \mathcal{C}\}$ are the weights. Fix an affine equation $l_H = 0$ for each hyperplane $H \in \mathcal{C}$. Set $Y = W - \bigcup_{H \in \mathcal{C}} H$. Consider the trivial line bundle $\mathcal{L}(a)$ over Y with an integrable connection $d(a) : \mathcal{O} \rightarrow \Omega^1$ given by $d + \Omega(a) = d + \sum_{H \in \mathcal{C}} a(H) d \log l_H$, where d is the de Rham differential. Denote by $\Omega^\bullet(\mathcal{L}(a))$ the complex of Y -sections of the homomorphic de Rham complex of $\mathcal{L}(a)$.

To any $H \in \mathcal{C}$ assign the one-form $i(H) = d \log l_H \in \Omega^1(\mathcal{L}(a))$. This construction defines a monomorphism $i(a) : (\mathcal{A}^\bullet(\mathcal{C}), d(a)) \rightarrow (\Omega^\bullet(\mathcal{L}(a)), d(a))$. The image of this monomorphism is called the *complex of the hypergeometric differential forms of weight a* . It is denoted by $(\mathcal{A}^\bullet(\mathcal{C}, a), d(a))$. The image of the homomorphism $i(a)S : (Fl^\bullet(\mathcal{C}), d) \rightarrow (\Omega^\bullet(\mathcal{L}(a)), d(a))$ is called the *complex of the flag hypergeometric differential forms of weight a* . It is denoted by $(Fl^\bullet(\mathcal{C}, a), d(a))$. For further details see [SV].

5.2 Discriminantal configurations

Let W be an affine complex space of dimension m . Let z_1, \dots, z_n be pairwise distinct complex numbers. Denote by \mathcal{C}_m a configuration in W consisting of hyperplanes $H_{kl} : t_k - t_l = 0$;

$1 \leq k < l \leq m$. So $\mathcal{C}_1 = \emptyset$, and $Y(\mathcal{C}_m)$ is the space of m -tuples of ordered distinct points in \mathbb{C} . Denote by $\mathcal{C}_{n;m}(z)$ a configuration in W consisting of hyperplanes $H_k^j : t_k - z_j = 0$, $1 \leq k \leq m$, $1 \leq j \leq n$, and H_{kl} , $1 \leq k < l \leq m$. Thus, $Y(\mathcal{C}_{n;m}(z)) = p^{-1}(z)$ where $p : Y(\mathcal{C}_{n+m}) \rightarrow Y(\mathcal{C}_n)$ is the projection on the first n coordinates. Define $\mathcal{C}_{0;m} = \mathcal{C}_m$.

Edges and flags of $\mathcal{C}_{n,m}$. For every non-empty subset $J = \{j_1, \dots, j_k\} \subset [m]$ set $L_J = H_{j_1 j_2} \cap H_{j_2 j_3} \cap \dots \cap H_{j_{k-1} j_k} \in \mathcal{C}_{n;m}^{k-1}$. L is an edge of codimension $k - 1$. In particular set $L_J = W$, for $k = 1$. For $i \in [n]$ define $L_J^i = H_{j_1}^i \cap H_{j_2}^i \cap \dots \cap H_{j_k}^i \in \mathcal{C}_{n;m}^k$. L_J^i is an edge of codimension k . Set $L_\emptyset^i = W$. Given non-intersecting subsets $J_1, \dots, J_k; I_1, \dots, I_n \subset [m]$, define $L_{J_1, \dots, J_k; I_1, \dots, I_n} = (\bigcap_{j=1}^k L_{J_j}) \cap (\bigcap_{i=1}^n L_{I_i}^i)$.

Multiplication of flags. Given two subsets $J \subset [m]$ and $I \subset [n]$, denote by $\mathcal{C}_{J;I} \subset \mathcal{C}_{m;n}$ the subset consisting of all hyperplanes $H_{j_1 j_2}$ with $j_1, j_2 \in J$ and H_j^i with $j \in J, i \in I$. Given subsets $J, J' \subset [m]$; $I, I' \subset [n]$ such that $J \cap J' = \emptyset$; $I \cap I' = \emptyset$, define maps $\circ : \text{Flag}^k(\mathcal{C}_{I;J}) \times \text{Flag}^l(\mathcal{C}_{I';J'}) \rightarrow \text{Flag}^{k+l}(\mathcal{C}_{I \cup I'; J \cup J'})$ as follows. For $F = F(H_1, \dots, H_k) \in \text{Flag}^k(\mathcal{C}_{I;J})$, $F' = F(H'_1, \dots, H'_l) \in \text{Flag}^l(\mathcal{C}_{I';J'})$, set $F \circ F' = (H_1, \dots, H_k, H'_1, \dots, H'_l)$. The following lemma, [SV] Lemma 5.7.2, takes place .

Lemma 5.2 *The above map correctly defines the map $Fl^k(\mathcal{C}_{I;J}) \otimes Fl^l(\mathcal{C}_{I';J'}) \rightarrow Fl^{k+l}(\mathcal{C}_{I \cup I'; J \cup J'})$. Moreover, for all $x \in Fl^k(\mathcal{C}_{I;J})$, $y \in Fl^l(\mathcal{C}_{I';J'})$ we have $x \circ y = (-1)^{kl} y \circ x$.*

5.3 Two maps of complexes

Let \mathfrak{g} be a Kac-Moody Lie algebra without Serre's relations. Let $M = M(\Lambda_1) \otimes \dots \otimes M(\Lambda_n)$ be a tensor product of Verma modules with weights $\Lambda_1, \dots, \Lambda_n \in \mathfrak{h}^*$. Set $\lambda = \underbrace{(1, 1, \dots, 1)}_m$.

In this case the number of generators $(f_j)_1^r$ of \mathfrak{n}_- equals m , i.e. $r = m$. Two maps of complexes ψ_\bullet and η_\bullet are described in [SV]:

$$\psi_p : C_p(\mathfrak{n}_-, M)_\lambda \rightarrow Fl^{m-p}, \quad \eta_p = \varphi^{-1} \circ (\psi_p^*)^{-1} : C_p(\mathfrak{n}_-, M^*)_\lambda \rightarrow A^{m-p}, \quad (43)$$

where φ is the map (40).

Note. The maps ψ_p define isomorphism of complexes. In view of Lemma 5.1 and [SV], Theorem 6.6 the maps $(-1)^{(m-p)(m-p-1)/2} \eta_p$ define isomorphism of complexes.

We will recall the explicit description of ψ_\bullet under the above assumption on λ . Let $g \in \mathfrak{n}_-$. A length $l = l(g)$ of a commutator g is given via an inductive definition. Set $l(f_j) = 1$ for $j = 1, \dots, m$. If $g = [g_1, g_2]$ and $l_1 = l(g_1)$, $l_2 = l(g_2)$, then set $l(g) = l_1 + l_2$. So $l(g) =$ "the number of f 's in g ".

To every commutator g assign a *bracket sign* $b(g) \in \mathbb{Z}/2\mathbb{Z}$ as follows. Set $b(f_j) = 0$; $b([g_1, g_2]) = b(g_1) + b(g_2) + l(g_1) \pmod{2}$.

To every commutator g assign a flag $Fl(g) \in Fl^{l(g)-1}(\mathcal{C}_{0;|g|})$ as follows. Set $Fl(f_j) = \square$. If $g = [g_2, g_1]$, set $Fl(g)$ equal to $Fl(g_1) \circ Fl(g_2)$ completed by the edge $L_{|g|}$.

Finally, for a commutator g set $F(g) = (-1)^{b(g)} Fl(g) \in Fl^{l(g)-1}(\mathcal{C}_{0;|g|})$. For $I = (i_1, \dots, i_l) \subset \{1, \dots, m\}$ and $1 \leq i \leq n$, set $f_I = f_{i_l} \dots f_{i_1} \in U(\mathfrak{n}_-)$ and $F^i(f_I) = F(H_{i_1}^i, \dots, H_{i_l}^i) \in Fl^l(\mathcal{C}_{\{i\};I})$. Let $z \in C_p(\mathfrak{n}_-, U(\mathfrak{n}_-)^{\otimes n})_\lambda$ and $z = g_p \wedge g_{p-1} \wedge \dots \wedge g_1 \otimes f_{I_n} \otimes f_{I_{n-1}} \otimes \dots \otimes f_{I_1}$,

where all g_i are commutators, $l_i = l(g_i)$. Let $\{f_{i_1}, \dots, f_{i_m}\}$ be the list of f_i 's in z read from right to left. Define $\sigma(z) \in \Sigma_m$ by $\sigma(z)(j) = i_j$. Set

$$\psi_p(z) = (-1)^{|\sigma(z)| + \sum_{i=1}^p (i-1)(l_i-1)} F^1(f_{I_1}) \circ \dots \circ F^n(f_{I_n}) \circ F(g_1) \circ \dots \circ F(g_p). \quad (44)$$

Note. There is a correction of the sign in the definition of ψ compared with [SV].

Examples. Let $n = 1$. $\psi(f_m \dots f_1) = F(H_1^1, \dots, H_m^1)$, and $\eta((f_{\sigma_1} \dots f_{\sigma_m})^*) = (-1)^{|\sigma|} H_{\sigma_1, \sigma_2} \circ \dots \circ H_{\sigma_{m-1}, \sigma_m} \circ H_{\sigma_m}^1$. Compose the inclusion map $i(a) : (\mathcal{A}^\bullet(\mathcal{C}), d(a)) \rightarrow (\Omega^\bullet(\mathcal{L}(a)), d(a))$ (see Section 5.1) with the map η to get

$$i(a) \circ \eta((f_{\sigma_1} \dots f_{\sigma_m})^*) = (-1)^{|\sigma|} d \ln(t_{\sigma_1} - t_{\sigma_2}) \wedge \dots \wedge d \ln(t_{\sigma_{m-1}} - t_{\sigma_m}) \wedge d \ln(t_{\sigma_m} - z_1).$$

Let $I \in P(\lambda, n)$ and $I = (i_1^1, \dots, i_{s_1}^1, \dots, i_1^n, \dots, i_{s_n}^n)$. Since $\lambda = (1, 1, \dots, 1)$, $I \in \Sigma_m$. Let $I_j = (i_{s_j}^j, \dots, i_1^j)$ for $1 \leq j \leq n$. We have $i(a) \circ \eta((f_{I_n})^* \otimes \dots \otimes (f_{I_1})^*) = (-1)^{|I|} \omega_I$. Therefore $i(a) \circ \eta(\sum_{I \in P(\lambda, n)} (f_I)^* f_I) = \omega(z, t)$, see Section 3.

Let $I = (i_1^1, \dots, i_{s_1}^1, \dots, i_1^n, \dots, i_{s_n}^n) \in P(\lambda, n)$, and $1 \leq k \leq s_j$. Define

$$\begin{aligned} f_{I; i_k^j} &= f_{I_n} \otimes \dots \otimes f_{I_{j+1}} \otimes f_{i_{k+1}^j} \dots f_{i_s^j} \otimes f_{I_{j-1}} \otimes \dots \otimes f_{I_1}, \\ \theta_{I; i_k^j} &= \omega_{i_1^1, \dots, i_{s_1}^1} \wedge \dots \wedge \omega_{i_1^{j-1}, \dots, i_{s_{j-1}}^{j-1}} \wedge \omega_{i_{k+1}^j, \dots, i_{s_j}^j} \wedge \omega_{i_1^{j+1}, \dots, i_{s_{j+1}}^{j+1}} \wedge \dots \wedge \omega_{i_1^n, \dots, i_{s_n}^n} \wedge \\ &\quad \wedge [d \ln(t_{i_1^j} - t_{i_2^j}) \wedge \dots \wedge d \ln(t_{i_{k-1}^j} - t_{i_k^j})] \\ H_{I; i_k^j} &= H_{i_1^1, \dots, i_{s_1}^1}^1 \circ \dots \circ H_{i_1^{j-1}, \dots, i_{s_{j-1}}^{j-1}}^{j-1} \circ H_{i_{k+1}^j, \dots, i_{s_j}^j}^j \circ H_{i_1^{j+1}, \dots, i_{s_{j+1}}^{j+1}}^{j+1} \circ \dots \circ H_{i_1^n, \dots, i_{s_n}^n}^n \circ \\ &\quad \circ [H_{i_1^j, i_2^j}^j \circ \dots \circ H_{i_{k-1}^j, i_k^j}^j], \end{aligned} \quad (45)$$

where $H_{i_1, \dots, i_l}^p = H_{i_1, i_2} \circ \dots \circ H_{i_{l-1}, i_l} \circ H_{i_l}^p$.

Lemma 5.3 *Let $I \in P(\lambda, n)$. Let $\epsilon_k^j = k((s_j - k) + s_{j+1} + \dots + s_n)$. Then*

$$i(a) \circ \eta(\Delta_{\sigma_{i_1^j}, \dots, \sigma_{i_k^j}} \otimes (f_{I; i_k^j})^*) = (-1)^{|I| + \epsilon_k^j} \theta_{I; i_k^j} \quad (46)$$

Proof. The statement of the Lemma is equivalent to the equation:

$$\Delta_{\sigma_{i_1^j}, \dots, \sigma_{i_k^j}} \otimes (f_{I; i_k^j})^* = (-1)^{|I| + \epsilon_k^j} \eta^{-1}(H_{I; i_k^j}). \quad (47)$$

It is sufficient to compute the two sides on elements of type $g \otimes f_{I; i_k^j}$ where g is a commutator of length k on $f_{i_1^j}, \dots, f_{i_k^j}$. Let $\sigma \in \Sigma_k$ and $f_{i_{\sigma_1}^j}, \dots, f_{i_{\sigma_k}^j}$ be the list of f_{i_j} 's entering g from right to left. The left hand side and the right hand side of (47) evaluated on $g \otimes f_{I; i_k^j}$ give

$$\begin{aligned} \Delta_{\sigma_{i_1^j}, \dots, \sigma_{i_k^j}} \otimes (f_{I; i_k^j})^* (g \otimes f_{I; i_k^j}) &= \Delta_{\sigma_{i_1^j}, \dots, \sigma_{i_k^j}}(g), \\ (-1)^{|I| + \epsilon_k^j} \eta^{-1}(H_{I; i_k^j})(g \otimes f_{I; i_k^j}) &= (-1)^{|I| + \epsilon_k^j} \varphi(H_{I; i_k^j})(\psi(g \otimes f_{I; i_k^j})), \end{aligned} \quad (48)$$

respectively. See formula (43). Use the definition of ψ to obtain

$$\begin{aligned} \psi(g \otimes f_{I;i_k^j}) &= (-1)^{|\tau|} F(H_{i_{s_1}^1}, \dots, H_{i_1^1}) \circ \dots \circ F(H_{i_{s_j}^j}, \dots, H_{i_{k+1}^j}) \circ \dots \\ &\quad \dots \circ F(H_{i_{s_n}^n}, \dots, H_{i_1^n}) \circ F(g), \end{aligned}$$

where $\tau = \left(\begin{smallmatrix} 1 & \dots & 1 & \dots & i_{s_j}^j & \dots & i_{k+1}^j & \dots & i_{s_n}^n & \dots & i_1^n & i_{\sigma_1}^j & \dots & i_{\sigma_k}^j \\ i_1^1 & \dots & i_{s_1}^1 & \dots & i_1^j & \dots & i_1^n & \dots & i_{s_n}^n & \dots & i_1^n & i_{\sigma_1}^j & \dots & i_{\sigma_k}^j \end{smallmatrix} \right)$. I as an element of Σ_m has the form $I = \left(\begin{smallmatrix} 1 & \dots & 1 & \dots & i_{s_j}^j & \dots & i_1^n & \dots & i_{s_n}^n \\ i_1^1 & \dots & i_{s_1}^1 & \dots & i_1^j & \dots & i_1^n & \dots & i_{s_n}^n \end{smallmatrix} \right)$. Thus

$$|I| = (|\tau| + S_k^j + k((s_j - k) + s_{j+1} + \dots + s_n) + |\sigma|) \pmod{2}, \quad (49)$$

where $S_k^j = \sum_{l=1, l \neq j}^n \frac{s_l(s_l - 1)}{2} + \frac{(s_j - k)(s_j - k - 1)}{2}$. Note that $H_{i_1, \dots, i_l}^p = (-1)^{l(l-1)/2} H_{i_l}^p \circ H_{i_{l-1}, i_l} \circ \dots \circ H_{i_1, i_2}$. Use the definition of φ , (40), to compute

$$\begin{aligned} \varphi(H_{i_1, \dots, i_l}^p) &= (-1)^{l(l-1)/2} \varphi(H_{i_l}^p \circ H_{i_{l-1}, i_l} \circ \dots \circ H_{i_1, i_2}) \\ &= (-1)^{l(l-1)/2} \delta_{F(H_{i_l}^p, \dots, H_{i_1}^p)} + \text{other } \delta - \text{summands}. \end{aligned} \quad (50)$$

$$\begin{aligned} \varphi(H_{I, i_k^j}) &= (-1)^{S_k^j} \varphi(H_{i_{s_1}^1}^1 \circ H_{i_{s_1-1}^1, i_{s_1}^1} \circ \dots \circ H_{i_1^1, i_2^1} \circ \dots \circ H_{i_{s_j}^j}^j \circ H_{i_{s_j-1}^j, i_{s_j}^j} \circ \dots \circ H_{i_{k+1}^j, i_{k+2}^j} \circ \\ &\quad \dots \circ H_{i_{s_n}^n}^n \circ H_{i_{s_n-1}^n, i_{s_n}^n} \circ \dots \circ H_{i_1^n, i_2^n} \circ [H_{i_1^j, i_2^j} \circ \dots \circ H_{i_{k-1}^j, i_k^j}]). \end{aligned} \quad (51)$$

We use formulae (49,50,51) to simplify (48).

$$(-1)^{|I|+e_k^j} \eta^{-1}(H_{I, i_k^j})(g \otimes f_{I; i_k^j}) = (-1)^{|\sigma|} \varphi(H_{i_1^j, i_2^j} \circ \dots \circ H_{i_{k-1}^j, i_k^j})(F(g)) \quad (52)$$

The proof of Lemma 5.3 is finished modulo the following result. \square

Lemma 5.4 *Let $\eta_{\mathcal{I}}$, $\psi_{\mathcal{I}}$ be the combinatorial maps (43), (44) defined on the set of distinct indices \mathcal{I} , $\mathcal{I} = \{i_1, \dots, i_k\} \subset \{1, \dots, m\}$. Then $\eta_{\mathcal{I}}(\Delta_{i_1, \dots, i_k}) = H_{i_1, i_2} \circ \dots \circ H_{i_{k-1}, i_k}$, for any $k = 2, \dots, m$.*

Proof. Induction by k . For $k = 2$, $g = [f_{i_1}, f_{i_2}]$ forms a base of the commutators of length 2 on f_{i_1} and f_{i_2} . $\Delta_{i_1, i_2}(g) = 1$. Since $b(g) = 1$ and $F(g) = (-1)^{b(g)} F(H_{i_1, i_2})$ and $\sigma = \begin{pmatrix} 1 & 2 \\ 2 & 1 \end{pmatrix}$, we have $\eta^{-1}(H_{i_1, i_2})([f_{i_1}, f_{i_2}]) = \varphi(H_{i_1, i_2})(\psi(g)) = \delta_{F(H_{i_1, i_2})}((-1)^{|\sigma|+b(g)} F(H_{i_1, i_2})) = 1$.

Let $2 < k \leq m$. Assume that for any j , $2 \leq j < k$, and $1 \leq s_1 < \dots < s_j \leq k$ we have $\eta(\Delta_{i_{s_1}, \dots, i_{s_j}}) = H_{i_{s_1}, i_{s_2}} \circ \dots \circ H_{i_{s_{j-1}}, i_{s_j}}$. Let g be a commutator of length k on f_{i_1}, \dots, f_{i_k} . Then $g = [g_1, g_2]$ with $l(g_1) = l_1$, and $l(g_2) = l_2$, and $l_1 + l_2 = k$. Let $\sigma \in \Sigma_k$ be such that $f_{i_{\sigma_1}}, \dots, f_{i_{\sigma_k}}$ is the list of f_i 's in g read from right to left. In order to evaluate $\eta^{-1}(H_{i_1, i_2} \circ \dots \circ H_{i_{k-1}, i_k})(g) = \varphi(H_{i_1, i_2} \circ \dots \circ H_{i_{k-1}, i_k})(\psi(g))$ remark that

$$\begin{aligned} \psi(g) &= (-1)^{|\sigma|+b(g)} (Fl(g_2) \circ Fl(g_1), L_{|g|}) = (-1)^{|\sigma|+l(g_1)} (F(g_2) \circ F(g_1), L_{i_1, \dots, i_k}) \\ \eta^{-1}(H_{i_1, i_2} \circ \dots \circ H_{i_{k-1}, i_k})(\psi(g)) &= \sum_{\tau \in \Sigma_{k-1}} (-1)^{|\tau|} \delta_{F(H_{i_{\tau_1}, i_{\tau_1+1}}, \dots, H_{i_{\tau_{k-1}}, i_{\tau_{k-1}+1}})}(\psi(g)) \end{aligned} \quad (53)$$

Since a link corresponding to a hyperplane $H_{j,j+1}$ connects only neighbouring indices in a flag of a type $F(H_{i_{\tau_1}, i_{\tau_1+1}}, \dots, H_{i_{\tau_{k-1}}, i_{\tau_{k-1}+1}})$ we have

$$\begin{aligned} & F(H_{i_{\tau_1}, i_{\tau_1+1}}, \dots, H_{i_{\tau_{k-1}}, i_{\tau_{k-1}+1}}) = \\ & = (\square, \dots, (t_{i_1} = \dots = t_{i_{\tau_{k-1}}}; t_{i_{\tau_{k-1}+1}} = \dots = t_{i_k}), L_{i_1, \dots, i_k} = (t_{i_1} = \dots = t_{i_k})). \end{aligned} \quad (54)$$

Let $\delta_{F(H_{i_{\tau_1}, i_{\tau_1+1}}, \dots, H_{i_{\tau_{k-1}}, i_{\tau_{k-1}+1}})}(\psi(g)) \neq 0$. Since $(-1)^{|\sigma|+b(g)}\psi(g) = (Fl(g_2) \circ Fl(g_1), L_{|g|}) = (\square, \dots, L_{|g_2|} \cap L_{|g_1|}, L_{|g|})$ formula (53) implies either $L_{|g_1|} = (t_{i_1} = \dots = t_{i_{\tau_{k-1}}}); L_{|g_2|} = (t_{i_{\tau_{k-1}+1}} = \dots = t_{i_k})$, or $L_{|g_2|} = (t_{i_1} = \dots = t_{i_{\tau_{k-1}}}); L_{|g_1|} = (t_{i_{\tau_{k-1}+1}} = \dots = t_{i_k})$. Without loss of generality we will assume that the second case takes place, i.e. $L_{|g_2|} = (t_{i_1} = t_{i_2} = \dots = t_{i_{\tau_{k-1}}}); L_{|g_1|} = (t_{i_{\tau_{k-1}+1}} = \dots = t_{i_{k-1}} = t_{i_k})$. Compare the lengths of the flags to conclude that $\tau_{k-1} = l_2$. In order to have non-zero multiples in the product

$$\begin{aligned} & \delta_{F(H_{i_{\tau_1}, i_{\tau_1+1}}, \dots, H_{i_{\tau_{k-1}}, i_{\tau_{k-1}+1}})}(Fl(g_2) \circ Fl(g_1), L_{|g|}) = \\ & = \delta_{F(H_{i_{\tau_1}, i_{\tau_1+1}}, \dots, H_{i_{\tau_{l_2-1}}, i_{\tau_{l_2-1}+1}})}(Fl(g_2))\delta_{F(H_{i_{\tau_{l_2}}, i_{\tau_{l_2}+1}}, \dots, H_{i_{\tau_{k-2}}, i_{\tau_{k-2}+1}})}(Fl(g_1)) \end{aligned} \quad (55)$$

we need $(\tau_1, \dots, \tau_{l_2-1})$ to be a permutation of the set $(1, \dots, l_2-1)$ and $(\tau_{l_2}, \dots, \tau_{k-2})$ to be a permutation of the set $(l_2+1, \dots, k-1)$. Set $\tau' = \begin{pmatrix} 1 & \dots & l_2-1 \\ \tau_1 & \dots & \tau_{l_2-1} \end{pmatrix}$ and $\tau'' = \begin{pmatrix} 1 & \dots & l_1-1 \\ \tau_{l_2-l_2} & \dots & \tau_{k-2-l_2} \end{pmatrix}$. Then $(-1)^{|\tau|} = (-1)^{|\tau'|+|\tau''|+l_1-1}$, $b(g) + l_1 = b(g_1) + b(g_2) \pmod{2}$, and

$$\begin{aligned} & \eta^{-1}(H_{i_1, i_2} \circ \dots \circ H_{i_{k-1}, i_k})(\psi(g)) = \\ & = (-1)^{|\sigma|+b(g)+l_1-1} \sum_{\tau' \in \Sigma_{l_2-1}} (-1)^{|\tau'|} \delta_{F(H_{i_{\tau'_1}, i_{\tau'_1+1}}, \dots, H_{i_{\tau'_{l_2-1}}, i_{\tau'_{l_2-1}+1}})}(Fl(g_2)) \times \\ & \times \sum_{\tau'' \in \Sigma_{l_1-1}} (-1)^{|\tau''|} \delta_{F(H_{i_{\tau''_{l_2}}, i_{\tau''_{l_2}+1}}, \dots, H_{i_{\tau''_{l_1-1}}, i_{\tau''_{l_1-1}+1}})}(Fl(g_1)) \\ & = (-1)^{|\sigma|-1} \eta^{-1}(H_{i_1, i_2} \circ \dots \circ H_{i_{l_2-1}, i_{l_2}})(F(g_2)) \eta^{-1}(H_{i_{l_2+1}, i_{l_2+2}} \circ \dots \circ H_{i_{k-1}, i_k})(F(g_1)) \\ & = (-1) \eta^{-1}(H_{i_1, i_2} \circ \dots \circ H_{i_{l_2-1}, i_{l_2}})(\psi(g_2)) \eta^{-1}(H_{i_{l_2+1}, i_{l_2+2}} \circ \dots \circ H_{i_{k-1}, i_k})(\psi(g_1)). \end{aligned} \quad (56)$$

The last equality holds because $\sigma = \begin{pmatrix} 1 & \dots & k \\ \sigma_1 & \dots & \sigma_k \end{pmatrix} = \begin{pmatrix} 1 & \dots & l_2 \\ \sigma_1 & \dots & \sigma_{l_2} \end{pmatrix} \times \begin{pmatrix} l_2+1 & \dots & l_k \\ \sigma_{l_2+1} & \dots & \sigma_{l_k} \end{pmatrix}$. Using the inductive hypothesis rewrite (56) as

$$\begin{aligned} \eta^{-1}(H_{i_1, i_2} \circ \dots \circ H_{i_{k-1}, i_k})(\psi(g)) & = (-1) \Delta_{i_1, \dots, i_{l_2}}(g_2) \Delta_{i_{l_2+1}, \dots, i_k}(g_1) \\ & = \Delta_{i_1, \dots, i_k}([g_1, g_2]). \end{aligned} \quad (57)$$

□

6 Derivation of the dynamical differential equation

In this section \mathfrak{g} will be a Kac-Mody Lie algebra without Serre's relations, $\lambda = (1, 1, \dots, 1)$, $r = m$. We will work in a weight space M_λ of the module $M = M(\Lambda_1) \otimes \dots \otimes M(\Lambda_n)$. We

will differentiate the hypergeometric form $\omega(z, t)$, express the result in terms of the complex $C_\bullet(\mathbf{n}_-, M^*)$, and derive the Dynamical differential equation in the form (25).

The integrand of a hypergeometric solution have the following form, see Section 3.

$$\begin{aligned}\Phi_\mu^{\frac{1}{\kappa}}\omega &= \exp\left(\frac{1}{\kappa}\left(-\sum_{i=1}^m \langle \alpha_{c(i)}, \mu \rangle t_i + \sum_{j=1}^n \langle \Lambda_j, \mu \rangle z_j\right)\right)\Phi_\mu^{\frac{1}{\kappa}}, \text{ where} \\ \Phi(z, t) &= \prod_{i < j} (z_i - z_j)^{\langle \Lambda_i, \Lambda_j \rangle} \prod_{k, j} (t_k - z_j)^{-\langle \alpha_{c(k)}, \Lambda_j \rangle} \prod_{k < l} (t_k - t_l)^{\langle \alpha_{c(k)}, \alpha_{c(l)} \rangle}.\end{aligned}$$

Fix $\mu' \in \mathfrak{h}$ and let $\partial_{\mu'}$ be the partial derivative with respect to the parameter μ in the direction of μ' . Then

$$\kappa \partial_{\mu'}(\Phi_\mu^{\frac{1}{\kappa}}\omega) = \left(-\sum_{i=1}^m \langle \alpha_{c(i)}, \mu' \rangle t_i + \sum_{j=1}^n \langle \Lambda_j, \mu' \rangle z_j\right)\Phi_\mu^{\frac{1}{\kappa}} \sum_{I \in P(\lambda, n)} (-1)^{|I|} \omega_I f_I v. \quad (58)$$

Let $I = (i_1^1, \dots, i_{s_1}^1; \dots; i_1^n, \dots, i_{s_n}^n) \in P(\lambda, n)$. Moreover $I \in \Sigma_m$ because of the form of λ . Since $r = m$ we have $c(i) = i$. Set $t_{i_{s_j+1}^j} = z_j$ and $\alpha_{i_{k;j}} = \alpha_{i_k^j} + \alpha_{i_{k-1}^j} + \dots + \alpha_{i_1^j}$. Rearrange the following expression:

$$\begin{aligned}\left(-\sum_{i=1}^m \langle \alpha_i, \mu' \rangle t_i + \sum_{j=1}^n \langle \Lambda_j, \mu' \rangle z_j\right)\omega_I &= \left(-\sum_{j=1}^n \sum_{k=1}^{s_j} \langle \alpha_{i_k^j}, \mu' \rangle t_{i_k^j} + \sum_{j=1}^n \langle \Lambda_j, \mu' \rangle z_j\right)\omega_I \\ &= \left(-\sum_{j=1}^n \sum_{k=1}^{s_j} \langle \alpha_{i_k^j}, \mu' \rangle (t_{i_k^j} - t_{i_{k+1}^j} + t_{i_{k+1}^j} - t_{i_{k+2}^j} + \dots + t_{i_{s_j}^j} - z_j + z_j) + \sum_{j=1}^n \langle \Lambda_j, \mu' \rangle z_j\right)\omega_I \\ &= \left(-\sum_{j=1}^n \sum_{k=1}^{s_j} \langle \alpha_{i_{k;j}}, \mu' \rangle (t_{i_k^j} - t_{i_{k+1}^j})\right)\omega_I + \left(\sum_{j=1}^n z_j \langle \Lambda_j - \alpha_{i_1^j} - \dots - \alpha_{i_{s_j}^j}, \mu' \rangle\right)\omega_I.\end{aligned} \quad (59)$$

Lemma 6.1 *Let $u(\mu, z) = \int_{\gamma(z)} \Phi_\mu^{\frac{1}{\kappa}}\omega$. Then*

$$\kappa \partial_{\mu'} u - \sum_{j=1}^n z_j \mu'^{(j)} u = \sum_{I \in P(\lambda, n)} (-1)^{|I|} \int_{\gamma(z)} \Phi_\mu^{\frac{1}{\kappa}} \left(-\sum_{j=1}^n \sum_{k=1}^{s_j} \langle \alpha_{i_{k;j}}, \mu' \rangle (t_{i_k^j} - t_{i_{k+1}^j})\right)\omega_I f_I v \quad (60)$$

Proof. Combine formulae (58), (59) with the fact $\mu'^{(j)} f_I v = \langle \Lambda_j - \alpha_{i_1^j} - \dots - \alpha_{i_{s_j}^j}, \mu' \rangle f_I v$ to obtain the result. \square

Lemma 6.2 *Define an operator L by $Lu = \kappa \partial_{\mu'} u - \sum_{j=1}^n z_j \mu'^{(j)} u$. Then*

$$Lu = \sum_{\alpha > 0} \frac{\langle \alpha, \mu' \rangle}{\langle \alpha, \mu \rangle} \Delta_{+, \alpha} u.$$

Proof. Let $I \in P(\lambda, n)$ and $1 \leq k \leq s_j$.

$$\begin{aligned}
& (t_{i_k^j} - t_{i_{k-1}^j})\omega_I = \omega_{i_1^1, \dots, i_{s_1}^1} \wedge \cdots \wedge (t_{i_k^j} - t_{i_{k-1}^j})\omega_{i_1^j, \dots, i_{s_j}^j} \wedge \cdots \wedge \omega_{i_1^n, \dots, i_{s_n}^n} \\
& = \omega_{i_1^1, \dots, i_{s_1}^1} \wedge \cdots \wedge [d \ln(t_{i_1^j} - t_{i_2^j}) \wedge \cdots \wedge d \ln(t_{i_{k-1}^j} - t_{i_k^j}) \wedge d(t_{i_k^j} - t_{i_{k+1}^j})] \wedge \\
& \wedge d \ln(t_{i_{k+1}^j} - t_{i_{k+2}^j}) \wedge \cdots \wedge d \ln(t_{i_{s_j}^j} - z_j)] \wedge \cdots \wedge \omega_{i_1^n, \dots, i_{s_n}^n} \\
& = (-1)^{\epsilon_k^j} \omega_{i_1^1, \dots, i_{s_1}^1} \wedge \cdots \wedge \omega_{i_{k+1}^j, \dots, i_{s_j}^j} \wedge \cdots \wedge \omega_{i_1^n, \dots, i_{s_n}^n} \wedge \\
& \wedge [d \ln(t_{i_1^j} - t_{i_2^j}) \wedge \cdots \wedge d \ln(t_{i_{k-1}^j} - t_{i_k^j})] \wedge d(t_{i_k^j} - t_{i_{k+1}^j}) \\
& = (-1)^{\epsilon_k^j} \theta_{I; i_k^j} \wedge d(t_{i_k^j} - t_{i_{k+1}^j}) = (-1)^{\epsilon_k^j + (m-1)} d(t_{i_k^j} - t_{i_{k+1}^j}) \wedge \theta_{I; i_k^j}, \tag{61}
\end{aligned}$$

where $\epsilon_k^j = k(s_{j+1} + \cdots + s_n + s_j - k)$. It is clear that $d\theta_{I; i_k^j} = 0$. Thus

$$\kappa d_t(\Phi_{\mu}^{\frac{1}{\kappa}} \theta_{I; i_k^j}) = \left(- \sum_{i=1}^m \langle \alpha_i, \mu \rangle dt_i + \Phi^{-1}(d_t \Phi)\right) \Phi_{\mu}^{\frac{1}{\kappa}} \theta_{I; i_k^j}. \tag{62}$$

Rearrange as in formula (59) and simplify to get

$$\begin{aligned}
& \left(\sum_{j=1}^n \sum_{k=1}^{s_j} \langle \alpha_{i_{k,j}}, \mu' \rangle d(t_{i_k^j} - t_{i_{k+1}^j})\right) \wedge \theta_{I; i_k^j} = \Phi^{-1}(d_t \Phi) \wedge \theta_{I; i_k^j} - \kappa \Phi_{\mu}^{-\frac{1}{\kappa}} d_t(\Phi_{\mu}^{\frac{1}{\kappa}} \theta_{I; i_k^j}) \\
& \langle \alpha_{i_{k,j}}, \mu' \rangle d(t_{i_k^j} - t_{i_{k+1}^j}) \wedge \theta_{I; i_k^j} = \Phi^{-1}(d_t \Phi) \wedge \theta_{I; i_k^j} - \kappa \Phi_{\mu}^{-\frac{1}{\kappa}} d_t(\Phi_{\mu}^{\frac{1}{\kappa}} \theta_{I; i_k^j}) \tag{63}
\end{aligned}$$

Since $\gamma(z)$ is a cycle, formulae (61), (63) allow us to rewrite Lu as

$$\begin{aligned}
Lu &= \sum_{I \in P(\lambda, n)} (-1)^{|I|} \int_{\gamma(z)} (-1)^{\epsilon_k^j} \left(- \sum_{j=1}^n \sum_{k=1}^{s_j} \frac{\langle \alpha_{i_{k,j}}, \mu' \rangle}{\langle \alpha_{i_{k,j}}, \mu \rangle} (\Phi_{\mu}^{\frac{1}{\kappa}} \Phi^{-1}(d_t \Phi)) \wedge \theta_{I; i_k^j} - \kappa d_t(\Phi_{\mu}^{\frac{1}{\kappa}} \theta_{I; i_k^j})\right) \\
&= \sum_{I \in P(\lambda, n)} \int_{\gamma(z)} \Phi_{\mu}^{\frac{1}{\kappa}} \left(- \sum_{j=1}^n \sum_{k=1}^{s_j} \frac{\langle \alpha_{i_{k,j}}, \mu' \rangle}{\langle \alpha_{i_{k,j}}, \mu \rangle} (-1)^{m-1} \frac{d_t \Phi}{\Phi} \wedge (i(a) \circ \eta(\Delta_{i_1^j, \dots, i_k^j} \otimes (f_{I; i_k^j})^*))\right) \tag{64}
\end{aligned}$$

Since $\theta_{I; i_k^j}$ is closed, its differential in the complex of the hypergeometric differential forms reduces to multiplication by $i(a)(\Omega(a)) = d_t \Phi / \Phi$. Taking into account Lemma 5.1 we have

$$\begin{aligned}
Lu &= \sum_{I \in P(\lambda, n)} f_I \int_{\gamma(z)} \Phi_{\mu}^{\frac{1}{\kappa}} \left(- \sum_{j=1}^n \sum_{k=1}^{s_j} \frac{\langle \alpha_{i_{k,j}}, \mu' \rangle}{\langle \alpha_{i_{k,j}}, \mu \rangle} (-1)^{m-1} d(a)(i(a) \circ \eta(\Delta_{i_1^j, \dots, i_k^j} \otimes (f_{I; i_k^j})^*))\right) \\
&= \sum_{I \in P(\lambda, n)} f_I \int_{\gamma(z)} \Phi_{\mu}^{\frac{1}{\kappa}} \left(- \sum_{j=1}^n \sum_{k=1}^{s_j} \frac{\langle \alpha_{i_{k,j}}, \mu' \rangle}{\langle \alpha_{i_{k,j}}, \mu \rangle} (i(a) \circ \eta(d\Delta_{i_1^j, \dots, i_k^j} \otimes (f_{I; i_k^j})^*))\right)
\end{aligned}$$

$$\begin{aligned}
&= \sum_{I \in P(\lambda, n)} f_I \int_{\gamma(z)} \Phi_{\mu}^{\frac{1}{\kappa}} \left(- \sum_{j=1}^n \sum_{k=1}^{s_j} \frac{\langle \alpha_{i_{k;j}}, \mu' \rangle}{\langle \alpha_{i_{k;j}}, \mu \rangle} (i(a) \circ \eta(\Delta_{i_1^j, \dots, i_k^j}(f_{I, i_k^j})^*)) \right) \\
&= \sum_{I \in P(\lambda, n)} f_I \int_{\gamma(z)} \Phi_{\mu}^{\frac{1}{\kappa}} \left(- \sum_{j=1}^n \sum_{k=1}^{s_j} \frac{\langle \alpha_{i_{k;j}}, \mu' \rangle}{\langle \alpha_{i_{k;j}}, \mu \rangle} (i(a) \circ \eta(\Delta_{i_1^j, \dots, i_k^j} e_{i_k^j}^j \dots e_{i_1^j}^j(f_I)^*)) \right) \quad (65)
\end{aligned}$$

Note that $\alpha_{i_{k;j}}$, $k = 1, \dots, s_j$ describe all λ -admissible roots such that $\Delta_{+, \alpha}^{(j)}(f_I)^* \neq 0$, see Section 4.4. Therefore

$$\begin{aligned}
Lu &= - \sum_{I \in P(\lambda, n)} f_I \int_{\gamma(z)} \Phi_{\mu}^{\frac{1}{\kappa}} \left(i(a) \circ \eta \left(\sum_{\alpha > 0} \frac{\langle \alpha, \mu' \rangle}{\langle \alpha, \mu \rangle} \sum_{(i_1, \dots, i_{m'}) \in P(\lambda_{\alpha}, 1)} \Delta_{i_1, \dots, i_{m'}} e_{i_{m'}} \dots e_{i_1}(f_I)^* \right) \right) \\
&= - \sum_{I \in P(\lambda, n)} f_I \int_{\gamma(z)} \Phi_{\mu}^{\frac{1}{\kappa}} \left(i(a) \circ \eta \left(\sum_{\alpha > 0} \frac{\langle \alpha, \mu' \rangle}{\langle \alpha, \mu \rangle} - \Delta_{-, \alpha}(f_I)^* \right) \right) \\
&= - \sum_{K, I \in P(\lambda, n)} f_I \int_{\gamma(z)} \Phi_{\mu}^{\frac{1}{\kappa}} \left(i(a) \circ \eta \left(\sum_{\alpha > 0} \frac{\langle \alpha, \mu' \rangle}{\langle \alpha, \mu \rangle} \langle -\Delta_{-, \alpha}(f_I)^*, f_K \rangle (f_K)^* \right) \right) \\
&= - \sum_{\alpha > 0} \frac{\langle \alpha, \mu' \rangle}{\langle \alpha, \mu \rangle} \sum_{K, I \in P(\lambda, n)} \langle -\Delta_{-, \alpha}(f_I)^*, f_K \rangle u_K f_I \quad (66)
\end{aligned}$$

$$Lu = - \sum_{\alpha > 0} \frac{\langle \alpha, \mu' \rangle}{\langle \alpha, \mu \rangle} \langle -\Delta_{-, \alpha}(f_I)^*, u \rangle f_I = - \sum_{\alpha > 0} \frac{\langle \alpha, \mu' \rangle}{\langle \alpha, \mu \rangle} \Delta_{+, \alpha} u \quad \square \quad (67)$$

The statement of Lemma 6.2 is equivalent to the Dynamical differential equation in the direction of μ' for a function u with values in the $(1, 1, \dots, 1)$ weight space of a \mathfrak{g} module M . The Symmetrization Lemma 4.11 deduces the general case from this one.

7 Main theorems

In this section we conclude the proofs of the the Theorems from Section 3 in the setting of Kac-Moody Lie algebras without Serre's relations. Then we deduce the corresponding results for any simple Lie algebra.

Let \mathfrak{g} be a Kac-Moody Lie algebra without Serre's relations. Let $\lambda \in \mathbb{N}^r$. Let $M = M(\Lambda_1) \otimes \dots \otimes M(\Lambda_n)$ be a tensor product of Verma modules for \mathfrak{g} with highest weights $\Lambda_1, \dots, \Lambda_n \in \mathfrak{h}^*$. Let $u(\mu, z) = \sum_{I \in P(\lambda, n)} u_I f_I$ be a hypergeometric integral with values in the weight space M_{λ} as described in Section 3, i.e. $u_I = \int_{\gamma(z)} \Phi_{\mu}^{\frac{1}{\kappa}} \left(\sum_{\sigma \in \Sigma(I)} (-1)^{|\sigma|} \omega_{I, \sigma} \right)$.

Theorem 7.1 *The function $u(\mu, z)$ solves the KZ equations (12) in M_{λ} .*

Proof. The proof given in Section 3 holds, because all relations we used are proved in [SV] in the general setting described above. \square

Theorem 7.2 *The function $u(\mu, z)$ solves the dynamical differential equations (13) in M_λ .*

Proof. Lemma 4.11 reduces the case of a general weight space M_λ to the case of a weight space $M_{\tilde{\lambda}}$, where $\tilde{\lambda} = \underbrace{(1, 1, \dots, 1)}_m$. Lemma 6.2 derives the Theorem in that case. \square

Proof of Theorem 3.2. Combine Corollary 4.4 and Theorem 7.2 to derive the dynamical differential equations for any Kac-Moody Lie algebra. In particular we have it for a simple Lie algebra. \square

Finally we will prove a determinant formula which establishes a basis of solutions for the system of KZ and dynamical differential equations in a weight space M_λ . From that formula we will derive the compatibility of the system of KZ and Dynamical differential equations.

Fix $\lambda \in \mathbb{N}^r$. Fix a basis $(f_I v)_{I \in P(\lambda, n)}$ of the weight space M_λ . Assume that a set $(\gamma_I(z))_{I \in P(\lambda, n)}$ of horizontal families of twisted cycles in $\{z\} \times \mathbb{C}^m$ is given

$$\text{Denote } u_{IJ} = \int_{\gamma_I(z)} \Phi_\mu^{\frac{1}{\kappa}} \left(\sum_{\sigma \in S(J)} (-1)^{|\sigma|} \omega_{J, \sigma} \right).$$

Proposition 7.3 *Let $\delta_\alpha = \text{tr}_{M_\lambda}(\Delta_{+, \alpha})$ for a positive root α of \mathfrak{g} . Denote $\epsilon_{ij} = \text{tr}_{M_\lambda}(\Omega_{ij, +})$. Then we have*

(a) *For any horizontal families of twisted cycles $(\gamma_I(z))_{I \in P(\lambda, n)}$ in $\{z\} \times \mathbb{C}^m$, there exists a constant $C = C(\Lambda_1, \dots, \Lambda_n, \lambda, \kappa)$ such that*

$$\det(u_{IJ}) = C \exp\left(\sum_{i=1}^n \frac{z_i}{\kappa} \text{tr}_{M_\lambda}(\mu^{(i)})\right) \prod_{\alpha > 0} \langle \alpha, \mu \rangle^{\frac{\delta_\alpha}{\kappa}} \prod_{i < j} (z_i - z_j)^{\frac{\epsilon_{ij}}{\kappa}}. \quad (68)$$

In the first product only finite number of factors are different from 1, i.e. $\delta_\alpha \neq 0$ if and only if $0 < \alpha \leq \lambda$.

(b) *For generic values of the parameters $(\Lambda_j)_{j=1}^m, (\alpha_i)_{i=1}^r, \kappa$ in a neighbourhood of a generic point $(\mu, z) \in \mathfrak{h} \times \mathbb{C}^n$ we can choose cycles $(\gamma_I(z))_{I \in P(\lambda, n)}$ such that the constant C from (a) is non-zero. Moreover the set of functions $\{u^I = \sum_{J \in P(\lambda, n)} u_{IJ} f_J\}_{I \in P(\lambda, n)}$ form a fundamental system of solutions for the system of KZ and dynamical differential equations.*

Proof. Part (a) is a corollary of Theorems 7.1 and 7.2. We will prove part (b) for values of the parameters such that all numbers $(\alpha_i, \alpha_j)/\kappa, -(\alpha_i, \Lambda_k)/\kappa$ have positive real parts for $1 \leq i, j \leq r, 1 \leq k \leq n$, and for a point (μ, z) such $z \in \mathbb{R}^n, z_1 < z_2 < \dots < z_n$ and $\langle \alpha_i, \mu \rangle / \kappa > 0$ for any $i = 1, \dots, r$. For generic values of $(\Lambda_j)_{j=1}^m, (\alpha_i)_{i=1}^r, z, \mu, \kappa$ (b) holds by analytic continuation.

The case $\lambda = \underbrace{(1, 1, \dots, 1)}_m$. Set $f_0(t) = \sum_{j=1}^m \langle \alpha_{c_\lambda(j)}, \mu \rangle t_j$. Let $I = (i_1^1, \dots, i_{s_1}^1; \dots; i_1^n, \dots, i_{s_n}^n) \in P(\lambda, n)$. Set $\gamma_I(z) = \{t \in \mathbb{R}^n : z_j < t_{i_1^j} < \dots < t_{i_{s_j}^j} < z_j + 1 \text{ for all } j = 1, \dots, n\}$, where $z_{n+1} = \infty$. Note that $\{\gamma_I(z)\}_{I \in P(\lambda, n)}$ is the set of all domains for the configuration of hyperplanes $H_{ij} : t_i - t_j = 0, H_i^k : t_i - z_k = 0, 1 \leq i < j \leq m, 1 \leq k \leq n$ which are either bounded, or the limit of f_0 on them is $+\infty$ when $\|t\| \rightarrow \infty$. In [Z] a linearly independent set of hypergeometric differential n -forms, called $\beta\mathbf{nbc}$ differential n -forms, associated

to those domains is defined. An explicit non-vanishing formula for the corresponding determinant is given in [MTV] Theorem 6.2, see also [DT]. Since $\lambda = (1, 1, \dots, 1)$ the space of hypergeometric n -forms is isomorphic to the space $C_0(\mathbf{n}_-^*, M^*)_\lambda$, see Section 5.3. The latter has basis $(f_I v^*)_{I \in P(\lambda, n)}$ which gives the basis $(\omega_I = i(a) \circ \eta(f_I v^*))_{I \in P(\lambda, n)}$ of the space of hypergeometric n -forms. Since this basis and the $\beta\mathbf{NBC}$ set have the same cardinality the non-zero determinant formula for the integrals of $\beta\mathbf{NBC}$ forms over the domains $\{\gamma_I(z)\}_{I \in P(\lambda, n)}$ implies a non-zero determinant formula for the integrals of $(\omega_I)_{I \in P(\lambda, n)}$ over the same domains. Since the determinant is non-zero at one point (μ, z) , it is non-zero at any point (μ, z) under the above conditions on the parameters.

The case of generic $\lambda \in \mathbb{N}^r$. Consider $C_0(\mathbf{n}_-^*, M^*)_\lambda$ and $C_0(\widetilde{\mathbf{n}}_-, \widetilde{M}^*)_{(1, 1, \dots, 1)}$ as in Section 4.4. A basis for the Σ_λ -symmetric hypergeometric differential forms is given by $(\omega_I)_{I \in P(\lambda, n)}$, where $\omega_I = \sum_{J \in \Sigma(I)} \widetilde{\omega}_J$ and $\widetilde{\omega}_J = i(a) \circ \eta(\widetilde{f}_J^*)$. $P(\underbrace{(1, 1, \dots, 1)}_m, n) = \cup_{I \in P(\lambda, n)} S(I)$

a disjoint union. Thus the set $(\omega_I)_{I \in P(\lambda, n)}$ consists of linearly independent forms in the space of all hypergeometric forms. The integral pairing described in the previous case is non-degenerate. Therefore there exists a subset of the set $(\gamma_J(z))_{J \in P((1, 1, \dots, 1), n)}$ indexed by the set $P(\lambda, n)$ such that the corresponding determinant is non-zero. \square

Corollary 7.4 *The system consisting of the union of KZ and Dynamic differential equations for any Kac-Moody Lie algebra with (or without) Serre's relations is a compatible system of differential equations.*

Remark. An algebraic proof of the compatibility of the system of KZ equations is given in [SV].

Proof. Let us write the differential operators which determine the KZ equations (12) and the dynamical equations (13) in the form

$$\text{KZ: } \frac{\partial}{\partial z_j} + B_j, \quad \text{Dynamical: } \frac{\partial}{\partial \mu'} + C_{\mu'}, \quad \text{where } j = 1, \dots, n \text{ and } \mu' \in \mathbf{h}. \quad (69)$$

The operators B_j and $C_{\mu'}$ are linear for any $j = 1, \dots, n$, $\mu' \in \mathbf{h}$. In order to prove the compatibility of the system of KZ and Dynamical differential equations we need to check $[\frac{\partial}{\partial z_j} + B_j, \frac{\partial}{\partial z_k} + B_k] = 0$, $[\frac{\partial}{\partial z_j} + B_j, \frac{\partial}{\partial \mu'} + C_{\mu'}] = 0$, and $[\frac{\partial}{\partial \mu'} + C_{\mu'}, \frac{\partial}{\partial \mu''} + C_{\mu''}] = 0$.

First consider the case of a Kac-Moody Lie algebra without Serre's relations, \mathfrak{g} , acting on a tensor product of highest weight modules M . We have

$$\left[\frac{\partial}{\partial z_j} + B_j, \frac{\partial}{\partial \mu'} + C_{\mu'} \right] = \left(\frac{\partial}{\partial z_j} C_{\mu'} \right) - \left(\frac{\partial}{\partial \mu'} B_j \right) + [B_j, C_{\mu'}] \quad (70)$$

The result is a linear operator with meromorphic coefficients depending on parameters $\{z, \mu, (\alpha_i)_{i=1}^r, (\Lambda_j)_{j=1}^n, \kappa\}$. Analogously, the commutators $[\frac{\partial}{\partial z_j} + B_j, \frac{\partial}{\partial z_k} + B_k]$ and $[\frac{\partial}{\partial \mu'} + C_{\mu'}, \frac{\partial}{\partial \mu''} + C_{\mu''}]$ are linear operators with meromorphic coefficients depending on the above set of parameters, where $j, k = 1, \dots, n$, $\mu', \mu'' \in \mathbf{h}$.

It is enough to show the commutativity of the above operators for generic values of the parameters. Then the commutators will be zero for any values of the parameters by analytic continuation.

Take such parameters $\{z, \mu, (\alpha_i)_{i=1}^r, (\Lambda_j)_{j=1}^n, \kappa\}$ that the set of hypergeometric solutions of the system of KZ and dynamical differential equations forms a basis of M . According to Proposition 7.3 (b) this is a generic choice of parameters. Since the KZ and the Dynamical differential operators act as zero on the set of hypergeometric solutions, their commutators also act as zero on the same set. Therefore the commutators act as zero on the \mathfrak{g} -module M .

Finally, consider a Kac-Moody Lie algebra with Serre's relations $\bar{\mathfrak{g}} = \mathfrak{g} / \ker(S : \mathfrak{g} \rightarrow \mathfrak{g}^*)$ which acts on $L = M / \ker(S : M \rightarrow M^*)$. Corollary 4.4 and [SV] Corollary 7.2.11 show that the Dynamical and the KZ operators for \mathfrak{g} correspond to the the the Dynamical and the KZ operators for $\bar{\mathfrak{g}}$ under this factorization. Then the commutativity of the operators on \mathfrak{g} implies that they are commutative on $\bar{\mathfrak{g}}$ as well. \square

References

- [BK] Babujian, H., Kitaev, A.: Generalized Knizhnik–Zamolodchikov equations and isomonodromy quantization of the equations integrable via the inverse scattering transform: Maxwell–Bloch system with pumping, *J. Math. Phys.* **39**, 2499–2506 (1998).
- [CVF] Chalykh, O. A., Feigin M. V., Veselov, A. P.: New integrable generalizations of Calogero–Moser quantum problem, *J. Math. Phys.* **39**, 695–703 (1998).
- [CV] Chalykh O. A., Veselov, A. P.: Commutative rings of partial differential operators and Lie algebras, *Commun. Math. Phys.* **126**, 597–611 (1990).
- [D] Drinfeld, V.: Quantum groups, *Proc. ICM (Berkeley, 1986)*, vol. 1, Amer. Math. Soc., 1987, 798–820.
- [DG] Duistermaat J. J., Grünbaum, F. A.: Differential operators in the spectral parameter, *Commun. Math. Phys.* **103**, 177–240 (1986).
- [DT] Douai, A., Terao, H.: The determinant of a hypergeometric period matrix, *Invent. Math.* **128**, 417–436 (1997).
- [HK] The bispectral problem (Montréal, 1997), Harnad, J., Kasman A.(eds.), CRM Proc. Lecture Notes **14**, Amer. Math. Soc., Providence, RI, 1998.
- [MTV] Markov, Y., Tarasov, V., Varchenko, A.: The determinant of a hypergeometric period matrix, *Houston J. Math.* **24**, vol.2, 197–219 (1998).
- [OS] Orlik, P., Solomon. L.: Combinatorics and topology of complements of hyperplanes, *Invent. Math.* **56**, 167–189 (1980).
- [SV] Schechtman, V., Varchenko, A.: Arrangements of hyperplanes and Lie algebra homology, *Inv. Math.* **106**, 139–194 (1991).
- [V] Varchenko, A.: Multidimensional hypergeometric functions and representation theory of Lie algebras and quantum groups, *Adv. Ser. Math. Phys.* **21**, World Scientific, 1995.
- [W] Wilson G., Bispectral commutative ordinary differential operators, *J. Reine Angew. Math.* **442**, 177–204 (1993).
- [Z] Ziegler, G.: Matroid shellability, β -systems, and affine arrangements, *J. Alg. Combinatorics*, **1**, 283–300 (1992).